

ONLINE SUPPLEMENTARY MATERIAL**Table S1:** Baseline Characteristics of excluded patients

	Excluded
Characteristics	n = 593
Male (n, %)	423
Age Group (years)	
< 65	242
65-74	171
≥ 75	180
Civil status (n, %)	
Married/Partnership	378
Divorced/Unmarried/Widow	>210
Missing	<5
Occupational status (n, %)	
Employed	158
Unemployed/out of workforce	417
Missing	18
Educational level (n, %)	
Low	256
Medium	200
High	52
Missing	72
Income, tertile (n, %)	
Low	245
Medium	197
High	>145
Missing	<5
Charlson Comorbidity Index	
Low (0 points)	421
Moderate/High (>0 points)	75
Missing	97

Table S2: Logistic regression model for those excluded compared with those included in the study population,

n = 1822

	n, (%)	Unadjusted		Multivariable adjusted*	
		OR	95% CI	OR	95% CI
Observations	593 (32.5)				
Civil status					
Married/Partnership	378 (20.7)	1 (ref.)		1 (ref.)	
Divorced/Unmarried/Widow	>210 (>11.5)	1.02	0.83-1.25	0.85	0.67-1.07
Missing	<5 (<1)				
Occupational status					
Employed	158 (8.7)	1 (ref.)		1 (ref.)	
Unemployed/Retired	417 (22.9)	1.69	1.36-2.09	1.54	1.14-2.08
Missing	18 (1.0)				
Educational level					
Low	256 (14.1)	1 (ref.)		1 (ref.)	
Medium	200 (10.1)	0.77	0.61-0.96	0.77	0.61-0.97
High	49 (2.7)	0.59	0.42-0.84	0.60	0.41-0.87
Missing	72 (4.0)				
Income, tertiles					
Low	245 (13.4)	1 (ref.)		1 (ref.)	
Medium	197 (10.8)	0.71	0.56-0.90	0.77	0.59-1.01
High	>145 (>7.9)	0.46	0.36-0.59	0.55	0.40-0.75
Missing	<5 (<0.3)				

* Adjusted for age, gender, ACS diagnosis, Charlson Comorbidity Index. OR: odds ratio; CI: confidence interval