

Table S2. List of fungal strains.

Name	Genotype / strain number	Origin
<u><i>Candida albicans</i> strains</u>		
KC2 = CAI4	<i>ura3Δ::imm434/ura3Δ::imm434</i>	(1)
KC68	KC2 <i>ccc2Δ::hisG/ccc2Δ::hisG</i>	(2)
KC139	KC68 <i>rbt5Δ::hisG/rbt5Δ::hisG</i>	(3)
KC485	KC68 <i>pga7Δ::hisG/pga7Δ::hisG</i>	(4)
KC782	KC68 <i>csa2Δ::hisG/csa2Δ::hisG</i>	(5)
<u>Other fungal species</u>		
<i>Candida parapsilosis</i>	Y2315	ARS NRRL ^a
<i>Candida auris</i>	Clinical isolate	Tel-Aviv Sourasky medical center – Ronen Ben-Ami
<i>Lodderomyces elongisporus</i>	CBS 2605	CBS ^b
<i>Meyerozima farinosa</i>	CBS 185	CBS
<i>Pichia stipitis</i>	CBS 5773	CBS

^a ARS NRRL = Agricultural Research Service culture collection, Peoria, IL

^b CBS = Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands.

SUPPLEMENTAL REFERENCES

1. Fonzi WA, Irwin MY. 1993. Isogenic strain construction and gene mapping in *Candida albicans*. *Genetics* 134:717–728.
2. Weissman Z, Shemer R, Kornitzer D. 2002. Deletion of the copper transporter CaCCC2 reveals two distinct pathways for iron acquisition in *Candida albicans*. *Mol Microbiol* 44:1551–1560.
3. Weissman Z, Kornitzer D. 2004. A family of *Candida* cell surface haem-binding proteins involved in haemin and haemoglobin-iron utilization. *Mol Microbiol* 53:1209–1220.
4. Kuznets G, Vigovsky E, Weissman Z, Lalli D, Gildor T, Kauffman SJ, Turano P, Becker J, Lewinson O, Kornitzer D. 2014. A Relay Network of Extracellular Heme-Binding Proteins Drives *C. albicans* Iron Acquisition from Hemoglobin. *PLoS Pathog* 10:e1004407.
5. Nasser L, Weissman Z, Pinsky M, Amartely H, Dvir H, Kornitzer D. 2016. Structural basis of haem-iron acquisition by fungal pathogens. *Nat Microbiol* 1:16156.