

Supplementary Table 1. *Podospora anserina* strains used in this research.

Strain	Genotype	Source
S+ (<i>Wt</i>)	<i>mat+</i>	Rizet, 1952
S- (<i>Wt</i>)	<i>mat-</i>	Rizet, 1952
<i>ku70</i>	Δ <i>ku70::geneticinR mat+</i>	R. Debuchy*
GFP-PTS1	<i>GFP-SKL::ble(EC) mat+</i>	Ruprich-Robert, et al, 2002
FS19	<i>FOX2::mCherry mat-</i>	Suaste-Olmos, et al, 2018
FS20	<i>FOX2::mCherry mat+</i>	Suaste-Olmos, et al, 2018
FS23	<i>FOX2::GFP mat+</i>	This study
FS24	<i>FOX2::GFP mat-</i>	This study
FS130	<i>IDH1::mCherry mat+</i>	This study
FS131	<i>IDH1::mCherry mat-</i>	This study
FS134	<i>ATP9-7::mCherry(EC) mat+</i>	This study
LP206	<i>FOX2::GFP ATP9-7::mCherry(EC) mat-</i>	This study
HT07	Δ <i>fis1::nat1 4D mat+</i>	This study
HT08	Δ <i>fis1::nat1 4D mat-</i>	This study
HT10	Δ <i>fis1::nat1 8B mat+</i>	This study
HT11	Δ <i>fis1::nat1 GFP-SKL::ble(EC) mat-</i>	This study
HT12	Δ <i>fis1::nat1 GFP-SKL::ble(EC) mat+</i>	This study
HT17	Δ <i>fis1::nat1 FOX2::mCherry GFP-SKL::ble(EC) mat-</i>	This study
HT18	Δ <i>fis1::nat1 FOX2::mCherry GFP-SKL::ble(EC) mat+</i>	This study
HT33	Δ <i>fis1::nat1 FOX2::mCherry mat+</i>	This study
HT34	Δ <i>fis1::nat1 FOX2::mCherry mat-</i>	This study
LP207	Δ <i>fis1::nat1 FOX2::GFP ATP9-7::mCherry(EC) mat+</i>	This study

HT35	$\Delta fisl1::nat1 FOX2::GFP ATP9-7::mCherry(EC) FIS1+(EC)-1 mat+$	This study
HT36	$\Delta fisl1::nat1 FOX2::GFP ATP9-7::mCherry(EC) FIS1+(EC)-6 mat+$	This study
FS132	$\Delta fisl1::nat1 FOX2::GFP mat+$	This study
FS133	$\Delta fisl1::nat1 FOX2::GFP mat-$	This study
RN1	$FOX2::GFP IDH1::mCherry mat+$	This study
RN5	$FOX2::GFP IDH1::mCherry mat-$	This study
RN10	$\Delta dnml1::hph FOX2::GFP mat+$	This study
RN11	$\Delta dnml1::hph FOX2::GFP mat-$	This study
RN13	$\Delta dnml1::hph IDH1::mCherry mat-$	This study
RN14	$\Delta dnml1::hph mat+$	This study
RN15	$\Delta dnml1::hph mat-$	This study
RN16	$\Delta dnml1::hph FOX2::GFP IDH1::mCherry mat+$	This study
RN17	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C1 mat+$	This study
RN23	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C5 mat+$	This study
RN18	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C9 mat+$	This study
RN19	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C11 mat+$	This study
RN20	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C12 mat+$	This study
RN21	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C13 mat+$	This study
RN22	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C14 mat+$	This study
RN24	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C21 mat+$	This study
RN25	$\Delta dnml1::hph FOX2::GFP DNM1+(EC)-C31 mat+$	This study

*Kindly provided by Dr. Robert Debuchy, Institute for Integrative Biology of the Cell, CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, France.

References

Suaste-Olmos, F., Zirion-Martinez, C., Takano-Rojas, H., and Peraza-Reyes, L. (2018). Meiotic development initiation in the fungus *Podospora anserina* requires the peroxisome receptor export machinery. *Biochim Biophys Acta* 1865(4), 572-586.

Rizet G. (1952). Les phénomènes de barrage chez *Podospora anserina*. I. Analyse génétique des barrages entre souches S and s. *Rev Cytol Biol Veg* 13, 51-92.

Ruprich-Robert, G., Berteaux-Lecellier, V., Zickler, D., Panvier-Adoutte, A., and Picard, M. (2002). Identification of six loci in which mutations partially restore peroxisome biogenesis and/or alleviate the metabolic defect of pex2 mutants in *Podospora*. *Genetics* 161(3), 1089-1099.