



Supplementary Materials

Supplementary Table 1. Medical conditions of the participants (n = 1,140).

Variable	Suicidality in Childhood						
	All	No (n = 1071)	Yes (n = 69)	p-Value			
Medical condition	n	n	n	%	n	%	
Hypertension	620	54.4	588	54.9	46.4	0.17	
Stroke	15	1.3	14	1.3	1	1.4	0.92
Heart diseases	116	10.2	101	9.4	15	21.7	0.001
Diabetes	172	15.1	160	14.9	12	17.4	0.58
Parkinson disease	5	0.4	5	0.5	0	0	0.57
Obesity (BMI \geq 30)	38	3.3	35	3.3	3	4.3	0.35
Psychopathology	9	0.8	8	0.7	1	1.4	0.52
Hyperlipidemia	189	16.6	173	16.2	16	23.2	0.13
Respiratory disease	48	4.2	45	4.2	3	4.3	0.95
Gastrointestinal disease	91	8	89	8.3	2	2.9	0.11
Urological disease	91	8	85	7.9	6	8.7	0.82
Musculoskeletal disease	110	9.6	100	9.3	10	14.5	0.16
Injury	15	1.3	12	1.1	3	4.3	0.02
Cancer	61	5.4	59	5.5	2	2.9	0.35
Hematological disease	9	0.8	7	0.7	2	2.9	0.04
Ophthalmological disease	201	17.6	181	16.9	20	29	0.01
Otorhinological disease	39	3.4	34	3.2	5	7.2	0.07
Sleep disorder	2	0.2	2	0.2	0	0	0.72
Other	97	8.5	90	8.4	7	10.1	0.62

Supplementary Table 2. Prevalence ratio with 95% CI for association with Geriatric Depression after MI ($n = 1,140$).

Variable.		Crude		Model I		Model II		Model III		Model IV	
		PR (95% CI)		APR (95% CI)							
<i>Childhood mental health status</i>											
Suicidal ideation (ref = No)	Yes	1.58	(1.15–2.15)	1.55	(1.17–2.07)	1.4	(1.04–1.88)	1.38	(1.03–1.86)	1.38	(1.02–1.88)
<i>Basic demographics</i>											
Age (years)		1.02	(1.01–1.03)	1.02	(1.01–1.03)	1.02	(1.01–1.03)	1.02	(1–1.03)	1.01	(0.99–1.02)
Sex (ref = Male)	Female	1.21	(0.99–1.47)	1.14	(0.94–1.38)	1.19	(0.98–1.45)	1.1	(0.9–1.35)	1.14	(0.91–1.43)
<i>Personality attribute</i>											
Neuroticism (ref = Q1)	Q2	1.49	(1.13–1.97)	1.29	(0.98–1.71)	1.29	(0.98–1.7)	1.26	(0.95–1.66)	1.22	(0.93–1.6)
Extroversion (ref = Q1)	Q3	2.01	(1.49–2.72)	1.61	(1.21–2.14)	1.57	(1.19–2.08)	1.5	(1.14–1.99)	1.39	(1.05–1.84)
Agreeableness (ref = Q1)	Q4	2.52	(1.93–3.29)	1.94	(1.48–2.53)	1.88	(1.44–2.45)	1.82	(1.39–2.37)	1.76	(1.36–2.28)
Conscientiousness (ref = Q1)	Q2	0.7	(0.55–0.89)	0.74	(0.58–0.94)	0.73	(0.58–0.93)	0.73	(0.58–0.93)	0.76	(0.59–0.96)
	Q3	0.56	(0.39–0.82)	0.66	(0.46–0.96)	0.65	(0.45–0.94)	0.67	(0.46–0.96)	0.67	(0.47–0.96)
	Q4	0.44	(0.33–0.59)	0.54	(0.4–0.73)	0.55	(0.45–0.74)	0.55	(0.41–0.75)	0.62	(0.46–0.83)
	Q2	0.65	(0.52–0.8)	0.68	(0.55–0.84)	0.67	(0.54–0.83)	0.66	(0.54–0.82)	0.68	(0.55–0.84)
	Q3	0.59	(0.41–0.84)	0.69	(0.48–0.97)	0.68	(0.48–0.96)	0.7	(0.49–0.98)	0.68	(0.49–0.95)
	Q4	0.29	(0.19–0.43)	0.41	(0.27–0.61)	0.41	(0.28–0.62)	0.42	(0.28–0.62)	0.43	(0.29–0.63)
	Q2	0.74	(0.57–0.95)	0.77	(0.6–0.98)	0.79	(0.62–1)	0.83	(0.65–1.06)	0.84	(0.66–1.06)
	Q3	0.73	(0.54–0.97)	0.71	(0.53–0.94)	0.72	(0.54–0.96)	0.78	(0.59–1.04)	0.81	(0.61–1.08)
	Q4	0.6	(0.43–0.84)	0.67	(0.48–0.93)	0.68	(0.48–0.95)	0.71	(0.51–0.99)	0.8	(0.57–1.11)

Openness (ref = Q1)	Q2	0.75	(0.56– –1)	0.9	(0.67 – 1.21)	0.94	(0.7– 1.26)	0.95	(0.71– 1.27)	0.93	(0.7– 1.24)
	Q3	0.66	(0.5– 0.87)	0.81	(0.63 – 1.06)	0.83	(0.64 – 1.08)	0.82	(0.64– 1.07)	0.85	(0.66– 1.09)
	Q4	0.49	(0.34– 0.69)	0.69	(0.48 – 0.97)	0.7	(0.49 – 0.99)	0.7	(0.5– 0.99)	0.72	(0.51– 1.01)
Childhood adversities											
ACE score (ref = 0)	1	1.67	(1.31– 2.14)		1.52	(1.2– 1.94)	1.45	(1.14– 1.84)	1.44	(1.13– 1.82)	
	2+	1.83	(1.4– 2.39)		1.53	(1.17– 2.01)	1.45	(1.11– 1.89)	1.35	(1.04– 1.76)	
Adulthood adversities											
Formal education (ref = <9 years)	10–12 years	0.82	(0.66– 1.02)			1.02	(0.82– 1.26)	1.05	(0.85– 1.31)		
	13 years +	0.6	(0.45– 0.82)			0.86	(0.64– 1.16)	0.91	(0.67– 1.23)		
Occupation (ref=manual/industrial laborer)	Skilled laborer	0.58	(0.41– 0.84)			0.75	(0.52– 1.07)	0.77	(0.53– 1.1)		
Annual household income (million yen) (ref = < 200)	200–500	0.6	(0.48– 0.75)			0.74	(0.59– 0.92)	0.79	(0.63– 1)		
	500–900	0.58	(0.41– 0.81)			0.67	(0.47– 0.94)	0.71	(0.49– 1.01)		
	900+	0.26	(0.10– 64)			0.33	(0.14– 0.8)	0.37	(0.15– 0.91)		
Total score of recent life event stressors	1	(1–1)				1	(1–1)	1	(1–1)		
Current health status & social relationships											
Total number of medical conditions	1.13	(1.05– 1.22)						1.07	(0.99– 1.16)		
Marital status (ref = Not married)	Married	0.65	(0.53– 0.79)					0.83	(0.65– 1.06)		
Living arrangement (ref = Alone)	With someone	0.69	(0.53– 0.9)					0.99	(0.73– 1.35)		
Employment status (ref = No)	Yes	0.7	(0.54– 0.89)					0.83	(0.65– 1.07)		
Giving support (ref = No)	Yes	0.61	(0.5– 0.75)					0.82	(0.63– 1.06)		
Receiving support (ref = No)	Yes	0.55	(0.46– 0.67)					0.78	(0.61– 1)		
Help-seeking attitude (ref = Passive)	Neutral	0.52	(0.41– 0.67)					0.63	(0.49– 0.79)		
	Active	0.59	(0.45– 0.78)					0.78	(0.6– 1.01)		

Note: MI = multiple imputation; APR = adjusted prevalence ratio; CI = confidential interval. Bold letters indicate statistically significant results where 95%CI did not include 1. Model I includes childhood suicidal ideation, demographics and personality attributes; Model II includes Model I variables + childhood adversities; Model III includes Model II variables + adulthood adversities; Model IV includes Model III variables + current health status and social relations.