

**Supplementary Table S1. Primers used in this study**

purpose for use	gene/vector	sequence
Vector construction of pENTR_mGFP	<i>mGFP</i>	CACCATGGTGAGCAAGGGCGAGGA TTACTTGTACAGCTCGTCATGCC
Vector construction of pENTR_LePGT1	<i>LePGT1</i>	GCAGGCTCCCGGGCCACCATGGTTCCAGCAAACAAACACAG AGCTGGGTCGGCGCAGCCAGGAACAACTTCCAAGTAAGATTCCG
Vector construction of pENTR_LePGT1	<i>pENTR/D-TOPO</i>	CGGCCGACCCAGCTTCTTGTACAAAGTT GGCCCGGAGGCTGCTTTTGACAAAGT
Vector construction of pGWB505_LeDI2	<i>LeDI2</i>	CACCATGGCTCAAGAAACTCTATTGCG TGGCAGCTCCATCCAGAAGG
Vector construction of pBin19_LeDI2-GFP	<i>LeDI2-GFP</i>	ATCTCTAGACCTTCACCATGGCTCAAGAAAACTC
Vector construction of pRI201AN_LeDI2-GFP	<i>LeDI2-GFP</i>	ATCGTCGACCCCTCACCATGGCTCAAGA
Vector construction of pBin19_LeDI2-GFP and pRI201AN_LeDI2-GFP	<i>LeDI2-GFP</i>	TTCGAGCTGGCCGCTTACTTGTAC
Site direct mutagenesis in <i>Xba</i> I in <i>LeDI2-GFP</i>	<i>LeDI2-GFP</i>	CATGGATCCTCTGATCGAACCACTTTGTACAA
Colony PCR and genotyping of hairy root	<i>GFP</i> <i>h</i>	GTGTTCGATCCAGAGGATCCATGGTGAGCAAG ATGGCCAGACTCACAAAGCATCATTG
Genotyping of hairy root	<i>Actin</i>	TCAAAGCTCATCGTGGTGGTGG GTGTTGGCTTATGAATTCCAGTC
Genotyping of hairy root	<i>GFP</i>	ATCTCAGATCCGGAAAGAAAGAAATACGA ATGGTGAGCAAGGGCGAGGAGCTGT
Colony PCR and genotyping of hairy root	<i>NPTII</i> (kanamycin resistant gene)	CTTGTACAGCTCGTCATGCCGTGAGTGAT ATGATTGAAACAAGATGGATTGC
Colony PCR and genotyping of hairy root	<i>HPT</i> (hygromycin resistant gene)	TCAGAAGAACCTCTCAAGAAGG TGCCCAGTCATAGCCGAATAGCCTCTCCAC CGGAAGTGCTTGACATTGG AGAAGAAGATGTTGGCGACC

**Supplementary Table S2. Medium composition**

	compound	amount (l <sup>-1</sup> )	callus induction medium	inoculation buffer	co-cultivation medium	root induction medium
1/2 MS	MgSO <sub>4</sub> · 7H <sub>2</sub> O	185 mg	+	+	+	+
	CaCl <sub>2</sub> · 2H <sub>2</sub> O	220 mg	+	+	-	+
	KNO <sub>3</sub>	950 mg	+	+	-	+
	NH <sub>4</sub> NO <sub>3</sub>	825 mg	+	+	-	+
	KH <sub>2</sub> PO <sub>4</sub>	85 mg	+	+	-	+
	FeSO <sub>4</sub> · 7H <sub>2</sub> O	13.9 mg	+	+	+	+
	Na <sub>2</sub> -EDTA	18.65 mg	+	+	+	+
	MnSO <sub>4</sub> · 4H <sub>2</sub> O	11.15 mg	+	+	+	+
	ZnSO <sub>4</sub> · 7H <sub>2</sub> O	4.3 mg	+	+	+	+
	CuSO <sub>4</sub> · 5H <sub>2</sub> O	0.0125 mg	+	+	+	+
	CoCl <sub>2</sub> · 6H <sub>2</sub> O	0.0125 mg	+	+	+	+
	KI	0.415 mg	+	+	+	+
	H <sub>3</sub> BO <sub>3</sub>	3.1 mg	+	+	+	+
	Na <sub>2</sub> MoO <sub>4</sub> · 2H <sub>2</sub> O	0.125 mg	+	+	+	+
	<i>myo</i> -inositol	100 mg	+	+	+	+
	Nicotinic acid	0.5 mg	+	+	+	+
	Pyridoxine hydrochloride	0.5 mg	+	+	+	+
	Thiamine hydrochloride	0.1 mg	+	+	+	+
	L-glycine	2 mg	+	+	+	+
	Sucrose	10 g	+	+	+	+
	Kinetin	100 µmol	+	-	-	-
	Indole acetic acid	10 µmol	+	-	-	-
	Acetosyringone	100 µmol	-	+	+	-
	Meropenem trihydrate	20 mg	-	-	-	+
	Agar	10 g	+	-	+	+

+, contained; -, not contained