

**Table S6.** Results From Sensitivity Meta-Analysis Including High-Risk-of-Bias Study

Endpoint	Efficacy Subgroup*	Within Subgroups				Between Subgroups (Heterogeneity)	
		No. of Studies	Pooled Hedges' <i>g</i> (95% CI)	<i>P</i>	<i>I</i> <sup>2</sup>	<i>Q</i> (df)	<i>P</i>
Total score	Non-efficacious	7	0.26 (0.01, 0.52)	0.043	0%	6.99 (1)	0.008
	Efficacious	10	0.67 (0.51, 0.83)	<0.001	74.1%		
Bowel symptoms	Non-efficacious	3	0.25 (-0.08, 0.58)	0.143	0%	3.53 (1)	0.060
	Efficacious	3	0.58 (0.47, 0.70)	<0.001	0%		
Systemic symptoms	Non-efficacious	3	0.14 (-0.18, 0.45)	0.271	0%	3.01 (1)	0.083
	Efficacious	3	0.41 (0.31, 0.52)	<0.001	0%		
Emotional function	Non-efficacious	3	0.24 (-0.09, 0.58)	0.148	28.2%	1.86 (1)	0.172
	Efficacious	3	0.49 (0.37, 0.60)	<0.001	0%		
Social function	Non-efficacious	3	0.08 (-0.25, 0.41)	0.626	0%	7.05 (1)	0.008
	Efficacious	3	0.55 (0.44, 0.67)	<0.001	0%		

\*Efficacious treatments are those which achieved primary endpoint; non-efficacious treatments are those for which primary endpoint was not met.