

## Supplementary Table S1

Table S1 Genital HSV-2 shedding and lesions rates using a Poisson regression method.

	Placebo N=28	Antigen only N=28	GEN-003, 10 µg N=31	GEN-003, 30 µg N=29	GEN-003, 100 µg N=27
Characteristic					
Rate Ratio for HSV shedding relative to pre-immunization (95% confidence interval)					
First period post-immunization	0.95 (0.57, 1.59)	1.29 (0.51, 3.27)	1.06 (0.57, 1.97)	0.47 (0.25, 0.85)*	0.68 (0.40, 1.14)
At 6 months	1.31 (0.78, 2.19)	1.61 (0.81, 3.17)	1.58 (0.94, 2.67)	0.55 (0.30, 1.03)	0.89 (0.57, 1.39)
Rate Ratio for lesions relative to pre-immunization (95% confidence interval)					
First period post-immunization	1.19 (0.55, 2.61)	0.64 (0.37, 1.13)	0.69 (0.33, 1.43)	0.53 (0.24, 1.16)	0.52 (0.23, 1.14)
At 6 months	1.38 (0.71, 2.71)	0.88 (0.43, 1.82)	0.89 (0.37, 2.14)	0.32 (0.15, 0.70)*	0.84 (0.39, 1.85)

\*p&lt;0.01

HSV, herpes simplex virus