Randomized controlled trial of medium cut-off versus high-flux dialyzers on quality of life outcomes in maintenance hemodialysis patients

Jeong-Hoon Lim¹, Yeongwoo Park², Ju-Min Yook¹, Soon-Youn Choi¹, Hee-Yeon Jung¹, Ji-Young Choi¹, Sun-Hee Park¹, Chan-Duck Kim¹, Yong-Lim Kim¹ and Jang-Hee Cho^{1*}

¹Department of Internal Medicine, School of Medicine, Kyungpook National University, Kyungpook National University Hospital, Daegu, South Korea. ²Department of Statistics, Kyungpook National University, Daegu, South Korea

Supplementary Information Listing:

Supplementary Table S1

*Corresponding author: Jang-Hee Cho, MD, PhD Associate Professor Division of Nephrology, Department of Internal Medicine Kyungpook National University Hospital, 130 Dongdeok-ro, Jung-gu, Daegu, 41944, Korea Tel: +82-53-200-5550 Fax: +82-53-426-2046

E-mail: jh-cho@knu.ac.k

Supplementary Table S1. Characteristics of dialyzers

Dialyzer	Membrane polymer	Membrane type	Inner diameter (μm)	Wall thickness (µm)	Effective surface area (m ²)	Ultrafiltration coefficient (mL/h/mmHg)	Priming volume (mL)	Sterilization
Therenova 400	Polyarylethersulfone- polyvinylpyrrolidone	Medium cut-off	180	35	1.7	48	91	Steam
FX CorDiax 80	Polysulfone- polyvinylpyrrolidone	High-flux	185	35	1.8	64	95	Steam
FX CorDiax 60	Polysulfone- polyvinylpyrrolidone	High-flux	185	35	1.4	47	74	Steam