

Electronic Supplementary Material

New markers for sepsis caused by *Pseudomonas aeruginosa* during burn infection

Moamen M. Elmassry¹, Nithya S. Mudaliar^{2*}, Jane A. Colmer-Hamood^{3,4}, Michael J. San Francisco^{1,5}, John A. Griswold², Sharmila Dissanaik², and Abdul N. Hamood^{2,3,†}

¹Department of Biological Sciences, Texas Tech University, Lubbock, TX, USA

²Department of Surgery, Texas Tech University Health Sciences Center, Lubbock, TX, USA

³Department of Immunology and Molecular Microbiology, Texas Tech University Health Sciences Center, Lubbock, TX, USA

⁴Department of Medical Education, Texas Tech University Health Sciences Center, Lubbock, TX, USA

⁵Honors College, Texas Tech University, Lubbock, TX, USA

*NM, current address: Caris Life Sciences, Phoenix, AZ, USA

†Corresponding author: Abdul N. Hamood, PhD; abdul.hamood@ttuhsc.edu

Online Resource 1: Table S1 Determination of *P. aeruginosa* systemic infection

Sample	Groups of Mice			
	Thermally injured and infected (B-I)	Thermally injured only (B-NI)	Infected only (NB-I)	Sham ^a (NB-NI)
Blood	~400 CFU/mL ^b	NG	NG	NG
Liver	125,333 ± 13,860 CFU/g ^c	NG	NG	NG
Spleen	103,000 ± 17,320 CFU/g ^c	NG	NG	NG

CFU, colony forming units; NG, no growth

^aMice were neither thermally injured nor infected.

^bTen µL of whole blood was plated to LB agar and *Pseudomonas* isolation agar; as the blood was clotting, the quantification is approximate.

^cLivers and spleens were individually weighed and homogenized in 1 mL PBS; the homogenates were serially diluted tenfold in PBS, and 10 µL of the dilutions plated on LB and *Pseudomonas* isolation agar to determine CFU present in the tissue. Values represent the means of three animals ± SEM.