

Supplementary Table 3. Birmingham IBS-S total scores and scores in the three domains of this scale in placebo and FMT-treated patients.

Time	Group	Total score	Pain	Diarrhoea	Constipation
0	Placebo	23.2±8.1	7.3±3.2	7.8±4.9	7.7±3.2
	30-g FMT	26.5±6.0	8.4±2.0	10.5±4.3**	7.5±3.4
	60-g FMT	25.2±6.8**	8.1±2.7	9.0±4.7	8.2±3.2
2 weeks	Placebo	19.6±10.6	6.2±4.2	6.0±6.0	7.4±3.6
	30-g FMT	20.4±8.1	7.3±3.3	6.8±4.5	5.5±3.6*
	60-g FMT	16.6±7.8	4.6±2.8***	5.9±4.7	6.2±4.1
1 month	Placebo	21.3±8.2	7.2±3.5	6.5±4.9	7.5±3.7
	30-g FMT	18.9±8.5	6.0±3.3	6.8±4.5	5.6±3.5*
	60-g FMT	15.8±8.6***	5.0±3.1**	4.9±4.0	6.0±4.8*
3 months	Placebo	19.8±9.7	6.6±3.9	6.2±5.3	7.0±4.5
	30-g FMT	15.3±7.0*	4.6±2.8**	4.9±3.0	5.7±3.8
	60-g FMT	14.6±7.2**	4.4±3.0**	4.2±2.9*	5.7±3.1

Data are mean±SD values.

*, p<0.05; **, p<0.01; ***, p<0.001 compared to placebo. ***, p<0.001 compared to 30-g FMT.