

Supplementary file: PICOS Search Strategy

PICOS	Population	No restriction
	Intervention	((prevent*[Title/Abstract]) AND (((intervention*[Title/Abstract]) OR program*[Title/Abstract]) OR strategy*) OR course[Title/Abstract]))
	Comparator	No restriction
	Outcome	((((((((depress*[Title/Abstract]) OR "depressive disorder"[Title/Abstract]) OR "depressive disorder"[MeSH Terms]) OR depression[Title/Abstract]) OR depression[MeSH Terms]) OR depression, postpartum[MeSH Terms])) AND (((((((((((postpartum[Title/Abstract]) OR postnatal[Title/Abstract]) OR puerperal[Title/Abstract]) OR perinatal[Title/Abstract]) OR prenatal[Title/Abstract]) OR antenatal[Title/Abstract]) OR intrapartum[Title/Abstract]) OR pregnancy[Title/Abstract]) OR pregnancy[MeSH Terms]) OR "pregnant women"[Title/Abstract]) OR "pregnant women"[MeSH Terms]) OR matern*[Title/Abstract]))))
	Study design	(((((("randomized controlled trial"[Publication Type]) OR random*[Title/Abstract]) OR controlled [Title/Abstract]) OR trial[Title/Abstract]) OR "clinical trial"[Title/Abstract]) OR "controlled clinical trial"[Publication Type])

No population and comparator restrictions included in the search strategy. The selection was performed using the inclusion and exclusion criteria.