

Stress disorders and psychiatric comorbidity

Supplemental Table 1. Number of Incident Psychiatric Diagnoses Following a Stress Disorder Diagnosis in Children (6 – 15 years) in Denmark, 1995-2011.

	Stress Cohort (N = 10,018)	Acute Stress Reaction (N = 854)	PTSD (N = 246)	Adjustment Disorder (N = 5,411)	ORSS (N = 600)	RSSU (N = 2907)	Comparison Cohort (N = 55,842)
Number of different incident psychiatric diagnoses (%)							
0	82	86	81	81	88	82	97
1	13	11	13	14	9.2	13	2.3
2+	4.4	3.0	5.3	4.4	3.3	4.8	0.5

Notes: PTSD = posttraumatic stress disorder. ORSS = Other reactions to severe stress. RSSU = Reactions to severe stress unspecified.

Stress disorders and psychiatric comorbidity

Supplemental Table 2. Average Time to Incident Psychiatric Diagnoses following a Stress Disorder Diagnosis in Children (6 – 15 years), in Denmark, 1995-2011.

Baseline stress disorder	Average time to incident psychiatric diagnoses (years)		
	Incident Depressive disorders	Incident Anxiety disorders	Incident Behavior disorders
Acute stress reaction	3.3	4.3	1.9
PTSD	3.8	4.2	3.4
Adjustment disorder	4.2	4.9	2.7
Other reactions to severe stress	3.7	3.0	3.2
Reactions to severe stress unspecified	3.5	4.1	2.1
Stress cohort	3.9	4.5	2.5
Comparison cohort	5.9	6.3	3.8