

**Genetically modified crops are superior in their nitrogen use efficiency-A meta-analysis of three major cereals**

Mengjiao Li<sup>1</sup>, Jili Xu<sup>1</sup>, Zhiyuan Gao<sup>1</sup>, Hui Tian<sup>1,\*</sup>, Yajun Gao<sup>1,\*</sup>& Khalil Kariman<sup>2</sup>

<sup>1</sup> Key Laboratory of Plant Nutrition and Agri-environment in Northwest China, Ministry of Agriculture, College of Natural Resources and Environment, Northwest A&F University, Yangling, Shaanxi, China

<sup>2</sup> School of Agriculture and Environment, The University of Western Australia, Crawley, WA 6009, Australia

Corresponding authors:

1. Hui Tian, tianh@nwsuaf.edu.cn
2. Yajun Gao, yajungao@nwafu.edu.cn

**Supplementary Table 7.** Gene categories involved in the collected studies

Gene name	Description	Category
<i>Bt</i>	Bt protein	
<i>Bar</i>	Herbicide resistance gene	<b>Biotic-stress related Protein (BSRP)</b>
<i>BjNPR1</i>	Pathogenesis-related genes 1	
<i>PcGDH</i>	Fungi NADP(H)-dependent GDH	
<i>ZmPti1-1</i>	Serine/threonine kinase The enzyme that mobilizes photoassimilated sucrose into	
<i>Incw2</i>	numerous reactions of the developing plant seeds, associated with grain yield	
<i>PDH45</i>	DEAD-box helicase	
<i>SAPK9</i>	Sucrose non-fermenting 1-related kinase	
<i>PEPC/PPDK</i>	Phosphoenolpyruvate carboxylase	
<i>appA</i>	Phytase	
<i>AVP1</i>	Vacuolar H <sup>+</sup> -pyrophosphatase	
<i>PLD</i>	Phospholipase D	
<i>VP</i>	V-H <sup>+</sup> Ppase	
<i>LRK1</i>	Leucine-rich repeat receptor-like kinase	
<i>AtGA2ox8</i>	Gibberellin 2-oxidase	
<i>SRK</i>	S-domain receptor-like kinase extracellular domain, involved in the self-incompatibility response	<b>Enzyme</b>
<i>AlaAt</i>	Alanine amino transferase	
<i>ASN</i>	Asparagine synthetase	
<i>GSI</i>	Cytosolic glutamine synthetase isoenzymes	
<i>ZmCIPK21</i>	CBL-Interacting Kinase	
<i>PsGPD</i>	Glyceral-Dehyde-3-phosphate dehydrogenase	
<i>SnRK</i>	Sucrose on-fermenting-1-related protein kinase	
<i>MDH</i>	Mitochondrial malate dehydrogenase	
<i>GA20-OXIDASE1</i>	GA biosynthesis	
<i>AVP1</i>	Arabidopsis vacuolar H <sup>+</sup> -pyrophosphatase	
<i>TPP</i>	Trehalose-6-phosphate phosphatase	
<i>TsVP</i>	Vacuolar H <sup>+</sup> -pyrophosphatase	
<i>BrECS</i>	Gamma-glutamylcysteine	

---

<i>LOSS</i>	synthetase
<i>Cu/Zn-SOD+APX</i>	Molybdenum cofactor sulfurase Superoxide dismutase/ascorbate peroxidase A vacuolar H <sup>+</sup>
<i>SaVHAC1</i>	-ATPase subunit c1 gene from the halophyte grass <i>Spartina alterniflora</i> Lo "isel
<i>SARK9</i>	Sucrose non-fermenting 1-related kinase 2 gene
<i>OsACS2</i>	1-aminocyclopropane-1-carboxylic acid synthase
<i>OsCPK4</i>	Calcium-dependent protein kinase
<i>OsHsFA2D</i>	Heat Shock transcription factor
<i>TsCBF1</i>	C-repeat-binding transcription factor
<i>NAC5, 6, 9, 67, 69</i>	NAC transcription factor
<i>AP37, 59</i>	Transcription factor
<i>ZmDOF1</i>	DOF transcription factor
<i>OsMYB55</i>	MYB transcription factor
<i>CCAAT Box-Binding Transcription Factor</i>	CCAAT Box-Binding transcription Factor
<i>OsPTF1</i>	Transcription factor
<i>DREB1A</i>	Encodes a member of the DREB subfamily A-1 of ERF/AP2 transcription factor family (CBF3)
<i>DHHC-type</i>	DHHC-type zinc finger protein
<i>Ossta2</i>	Oryza sativa salt tolerance activation 2-Dominant
<i>W16</i>	AP2/EREBP transcription factor
<i>DHHC</i>	Zinc-finger protein
<i>OsHAK1</i>	High-affinity potassium transporter
<i>NAT3</i>	A nitrate transporter of cylindrotheca fusiformis
<i>OsPTR6</i>	Nitrate transporter
<i>STP13</i>	Hexose transporter
<i>ApKUP3</i>	High-affinity potassium transporters
<i>PHO1</i>	Pi transporter
<i>PgNHX1</i>	Pennisetum glaucum vacuolar Na <sup>+</sup> /H <sup>+</sup> antiporter gene
	Transporter

---

---

<i>nhaA</i>	Na <sup>+</sup> /H <sup>+</sup> antiporter
<i>HvSUT1</i>	Sucrose transporter
<i>OsNRT2.3b/OsNPF7.3/OsNRT1.1A.OsNPF6.3/OsNR2.1/OsNPF7.9/OsNRT1.2</i>	Nitrate transporter
<i>OsAMT1-1</i>	Ammonium transporter
<i>HVA1</i>	Late embryogenesis abundant protein
<i>AISAP</i>	Stress associated protein
<i>OsEOD93-1</i>	Nitrogen-responsive early nodulin gene
<i>Tin</i>	Tiller inhibition gene
<i>OCP11</i>	Stress responsive proteinase inhibitor gene
<i>Bn-csRRM2</i>	A conserved RNA-binding motif (RRM) domain, flowering control locus A (FCA) protein
<i>TaPHR1</i>	A phosphate starvation response regulator
<i>LEA</i>	Late embryogenesis abundant
<i>SeCspA</i>	Synthetic bacterial cold shock protein gene
<i>TOND1</i>	Tolerance of nitrogen deficiency
<i>OsPIN2</i>	Efflux-facilitating PIN-FORMED
<i>TaMIR1129</i>	miRNA
<i>OsDhn1</i>	Dehydrin gene
<i>OsCam1-1</i>	Salt-inducible calmodulin gene
<i>RAA1</i>	Root architecture associated 1
<i>MoSMI</i>	Immunity-inducing protein
<i>SaARF1</i>	ADP ribosylation factor
<i>RPL23A</i>	Ribosomal protein large
<i>PgRab7</i>	Vesicle trafficking protein
<i>AtGRP2</i>	Glycine-rich RNA-binding protein
<i>SeCspA</i>	Synthetic bacterial cold shock protein
<i>U</i>	Unknown genes
<i>EhEM1</i>	Em-like protein from <i>Eutrema halophilum</i>
<i>OsENOD93-1</i>	A nitrogen-responsive early nodulin

---

---

*OsPGIPI* gene  
Polygalacturonase-inhibiting  
proteins

---