

Supplementary Table S2: Antifungal susceptibility comparison between *C. tropicalis* and *C. albicans* blood isolates

Antifungal drugs	Species	Category number (%)			P-value
		S/WT	SDD/I	R/NWT	
Voriconazole	<i>C. tropicalis</i>	21 (43.8)	6 (12.4)	21 (43.8)	< 0.001
	<i>C. albicans</i>	45 (97.8)	0 (0)	1 (2.2)	
Fluconazole	<i>C. tropicalis</i>	26 (54.1)	1 (2.1)	21 (43.8)	< 0.001
	<i>C. albicans</i>	45 (97.8)	0 (0)	1 (2.2)	
Posaconazole ^a	<i>C. tropicalis</i>	30 (62.5)	NA	18 (37.5)	< 0.001
	<i>C. albicans</i>	45 (97.8)	NA	1 (2.2)	
Anidulafungin	<i>C. tropicalis</i>	47 (97.9)	0 (0)	1 (2.1)	0.325
	<i>C. albicans</i>	46 (100)	0 (0)	0 (0)	
Micafungin	<i>C. tropicalis</i>	47 (97.9)	0 (0)	1 (2.1)	0.325
	<i>C. albicans</i>	46 (100)	0 (0)	0 (0)	
Caspofungin	<i>C. tropicalis</i>	47 (97.9)	0 (0)	1 (2.1)	0.325
	<i>C. albicans</i>	46 (100)	0 (0)	0 (0)	
Amphotericin B ^a	<i>C. tropicalis</i>	48 (100)	NA	0 (0)	NA
	<i>C. albicans</i>	46 (100)	NA	0 (0)	

Notes: ^acategorization based on the epidemiologic cut-off values (ECVs); Itraconazole and 5-flucytosine interpretation were not available for *C. albicans*. Abbreviations: MIC, minimal inhibitory concentration; GM, geometric mean; S, susceptible; WT, wild type; SDD, susceptible-dose dependent; I, intermediate; R, resistant; NWT, non-wild type; NA, not applicable