

Supplementary information

**Guidelines and definitions for research on
epithelial–mesenchymal transition**

In the format provided by the
authors and unedited

Supplementary Table:**Additional transcription factors associated with EMT and key studies reporting their discovery in the context of development, cancer and fibrosis.**

Transcription factor*	Type	Development	Cancer	Fibrosis
Alx1	Homeobox	Saunders & McClay, 2014	Yuan et al., 2013	-
AP1 (Jun/Fos Atf)	Ap1	Reichmann et al., 1992	Bakiri et al. 2015	Alcorn et al., 2009
Brachyury (T)	T-box	Beddington et al., 1992	Fernando et al., 2010	Sun et al., 2014
CBF-A (NF-YA)	CCAT box binding	-	Venkov et al., 2007	Venkov et al., 2007
E2-2 (TCF4)	bHLH	Sobrado et al., 2009	Sobrado et al., 2009	-
E47 (TCF3)	bHLH	Perez-Moreno et al., 2001	Perez-Moreno et al., 2001	Slattery et al., 2006
Erg	ETS	Saunders & McClay, 2014	Petrovics et al., 2005	-
Ets1	ETS	Fafeur et al., 1997	Nakayama et al., 1996	Nakamura et al., 2004.
FoxC2 (FoxB)	Forkhead	Saunders & McClay, 2014	Mani et al., 2007	Sipos & Galamb, 2012
GATA4/6 (Srp)	GATA	Campbell et al, 2011	-	-
Gooseoid (GSC)	Homeobox	Blum et al., 1992	Hartwell et al., 2006	-
HoxB7	Homeobox	-	Wu et al, 2006	-
Id1	HLH	Jen et al., 1996	Tobin et al, 2011	Li et al., 2007
KLF2	Sp/KLF	Chiplunkar et al., 2013	-	-
Klf4	Klf	Liu et al., 2012	Lin et al., 2012	Tiwari et al., 2013
KLF6 (Z9f)	Sp/KLF	-	-	Holian et al., 2008
KLF8	Sp/KLF	-	Wang et al, 2007	-
NF-κb	NF- κ B/Rel	DeLaughter et al., 2016	Huber et al, 2004	Sunami et al., 2012
Prrx1	Paired related homeobox	Ocana et al. 2012	Ocana et al., 2012; Takano et al., 2016	Gong et al., 2017
Runx2	Runt-related	Tavares et al. 2017	Pratap et al., 2005	Mummler et al., 2018

Six1	Homeobox	Grifone et al., 2005	Micalizzi et al., 2009; McCoy et al., 2009	Smith et al., 2008
Sox4	Homeobox	Zhang et al, 2012	Liu et al., 2006	Xiao et al., 2015
Sox9	Homeobox	Akiyama et al., 2004	Acevedo et al., 2007	Hanley et al., 2008
Tbx3	T-box	Singh et al., 2012	Rodriguez et al. 2008	Wensing & Campos, 2014
Tead (Yap/Taz)	Tead	Diepenbruck et al., 2014	Overholtzer et al., 2006	Seo et al 2016
Tgif1	Homeobox	Saunders & McClay, 2014	Xiang et al., 2015	-
Twist2 (Dermo1)	bHLH	-	Ansieau et al., 2008	Grunz-Borgmann et al., 2017

*Transcription factors with a minimum of 10 citations in the EMT literature. “-“ indicates lack of a pertinent reference.

References:

- Acevedo, V.D., Gangula, R. D., Freeman, K. W., Li, R. L., Zhang, Y. Y., Wang, F., Ayala, G. E., Peterson, L. E., Ittmann, M. & Spencer, D. M. Inducible FGFR-1 activation leads to irreversible prostate adenocarcinoma and an epithelial-to-mesenchymal transition. *Cancer Cell* 12, 559-571 (2007).
- Akiyama, H., Chaboissier, M. C., Behringer, R. R., Rowitch, D. H., Schedl, A., Epstein, J. A. & de Crombrughe, B. Essential role of Sox9 in the pathway that controls formation of cardiac valves and septa. *Proc Natl Acad Sci U S A* 101, 6502-6507 (2004).
- Alcorn, J.F., van der Velden, J., Brown, A. L., McElhinney, B., Irvin, C. G. & Janssen-Heininger, Y. M. W. c-Jun N-Terminal Kinase 1 Is Required for the Development of Pulmonary Fibrosis. *American Journal of Respiratory Cell and Molecular Biology* 40, 422-432 (2009).
- Ansieau, S., Bastid, J., Doreau, A., Morel, A. P., Bouchet, B. P., Thomas, C., Fauvet, F., Puisieux, I., Doglioni, C., Piccinin, S., Maestro, R., Voeltzel, T., Selmi, A., Valsesia-Wittmann, S., Caron de Fromentel, C. & Puisieux, A. Induction of EMT by twist proteins as a collateral effect of tumor-promoting inactivation of premature senescence. *Cancer Cell* 14, 79-89 (2008).
- Bakiri, L., Macho-Maschler, S., Custic, I., Niemiec, J., Guio-Carrion, A., Hasenfuss, S. C., Eger, A., Muller, M., Beug, H. & Wagner, E. F. Fra-1/AP-1 induces EMT in mammary epithelial cells by modulating Zeb1/2 and TGF beta expression. *Cell Death Differ* 22, 336-350 (2015).
- Beddington, R.S.P., Rashbass, P.R. & Wilson, V. Brachyury--a gene affecting mouse gastrulation and early organogenesis. *Development*.-Supplement., 157-165 (1992).

- Blum, M., Gaunt, S. J., Cho, K. W., Steinbeisser, H., Blumberg, B., Bittner, D., & De Robertis, E. M. Gastrulation in the mouse: the role of the homeobox gene goosecoid. *Cell* 69, 1097-1106 (1992).
- Campbell, K., Whissell, G., Franch-Marro, X., Batlle, E. & Casanova, J. Specific GATA factors act as conserved inducers of an endodermal-EMT. *Dev Cell* 21, 1051-1061, doi:10.1016/j.devcel.2011.10.005 (2011)
- Chiplunkar, A.R., Lung, T. K., Alhashem, Y., Koppenhaver, B. A., Salloum, F. N., Kukreja, R. C., Haar, J. L. & Lloyd, J. A. Kruppel-like factor 2 is required for normal mouse cardiac development. *Plos One* 8, e54891 (2013).
- DeLaughter, D.M., Clark, C. R., Christodoulou, D. C., Seidman, C. E., Baldwin, H. S., Seidman, J. G. & Barnett, J. V. Transcriptional Profiling of Cultured, Embryonic Epicardial Cells Identifies Novel Genes and Signaling Pathways Regulated by TGFbetaR3 In Vitro. *Plos One* 11, e0159710 (2016).
- Diepenbruck, M., Waldmeier, L., Ivanek, R., Berninger, P., Arnold, P., van Nimwegen, E. & Christofori, G. Tead2 expression levels control the subcellular distribution of Yap and Taz, zyxin expression and epithelial-mesenchymal transition. *Journal of Cell Science* 127, 1523-1536 (2014).
- Fafeur, V., Tulasne, D., Queva, C., Vercamer, C., Dimster, V., Mattot, V., Stehelin, D., Desbiens, X. & Vandembunder, B. The ETS1 transcription factor is expressed during epithelial-mesenchymal transitions in the chick embryo and is activated in scatter factor-stimulated MDCK epithelial cells. *Cell Growth & Differentiation* 8, 655-665 (1997).
- Fernando, R.I., Litzinger, M., Trono, P., Hamilton, D. H., Schlom, J. & Palena, C. The T-box transcription factor Brachyury promotes epithelial-mesenchymal transition in human tumor cells. *J Clin Invest* 120, 533-544 (2010).
- Gong, J., Han, J., He, J. Y., Liu, J. M., Han, P., Wang, Y. W., Li, M. K., Li, D. X., Ding, X. M., Du, Z. P., Liao, J. Z. & Tian, D. Paired related homeobox protein 1 regulates PDGF-induced chemotaxis of hepatic stellate cells in liver fibrosis. *Laboratory Investigation* 97, 1020-1032 (2017).
- Grifone, R., Demignon, J., Houbron, C., Souil, E., Niro, C., Seller, M. J., Hamard, G. & Maire, P. Six1 and Six4 homeoproteins are required for Pax3 and Mrf expression during myogenesis in the mouse embryo. *Development* 132, 2235-2249 (2005).
- Grunz-Borgmann, E.A., Nichols, L.A., Wang, X.H. & Parrish, A.R. Twist2 Is Upregulated in Early Stages of Repair Following Acute Kidney Injury. *Int J Mol Sci* 18 (2017).
- Hanley, K.P., Oakley, F., Sugden, S., Wilson, D. I., Mann, D. A. & Hanley, N. A. Ectopic SOX9 mediates extracellular matrix deposition characteristic of organ fibrosis. *Journal of Biological Chemistry* 283, 14063-14071 (2008).

- Hartwell, K.A., Muir, B., Reinhardt, F., Carpenter, A. E., Sgroi, D. C. & Weinberg, R. A. The Spemann organizer gene, Goosecoid, promotes tumor metastasis. *Proc Natl Acad Sci U S A* 103, 18969-18974 (2006).
- Holian, J., Qi, W., Kelly, D. J., Zhang, Y., nMreich, E., Pollock, C. A. & Chen, X. M. Role of Kruppel-like factor 6 in transforming growth factor-beta1-induced epithelial-mesenchymal transition of proximal tubule cells. *Am J Physiol Renal Physiol* 295, F1388-1396 (2008).
- Huber, M.A., Azoitei, N., Baumann, B., Grunert, S., Sommer, A., Pehamberger, H., Kraut, N., Beug, H. & Wirth, T. NF-kappa B is essential for epithelial-mesenchymal transition and metastasis in a model of breast cancer progression. *Journal of Clinical Investigation* 114, 569-581 (2004).
- Jen, Y., Manova, K. & Benezra, R. Expression patterns of Id1, Id2, and Id3 are highly related but distinct from that of Id4 during mouse embryogenesis. *Developmental Dynamics* 207, 235-252 (1996).
- Li, Y., Yang, J., Luo, J.H., Dedhar, S. & Liu, Y. Tubular epithelial cell dedifferentiation is driven by the helix-loop-helix transcriptional inhibitor Id1. *J Am Soc Nephrol* 18, 449-460 (2007).
- Lin, Z.S., Chu, H.C., Yen, Y.C., Lewis, B.C. & Chen, Y.W. Kruppel-like factor 4, a tumor suppressor in hepatocellular carcinoma cells reverts epithelial mesenchymal transition by suppressing slug expression. *Plos One* 7, e43593 (2012).
- Liu, P.B., Ramachandran, S., Ali Seyed, M., Scharer, C. D., Laycock, N., Dalton, W. B., Williams, H., Karanam, S., Datta, M. W., Jaye, D. L. & Moreno, C. S. Sex-determining region Y box 4 is a transforming oncogene in human prostate cancer cells. *Cancer Research* 66, 4011-4019 (2006).
- Liu, Y.N., Abou-Kheir, W., Yin, J. J., Fang, L., Hynes, P., Casey, O., Hu, D., Wan, Y., Seng, V., Sheppard-Tillman, H., Martin, P. & Kelly, K. Critical and reciprocal regulation of KLF4 and SLUG in transforming growth factor beta-initiated prostate cancer epithelial-mesenchymal transition. *Mol Cell Biol* 32, 941-953 (2012).
- Mani, S.A., Yang, J., Brooks, M., Schwaninger, G., Zhou, A., Miura, N., Kutok, J. L., Hartwell, K., Richardson, A. L. & Weinberg, R. A. Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and is associated with aggressive basal-like breast cancers. *Proceedings of the National Academy of Sciences of the United States of America* 104, 10069-10074 (2007).
- McCoy, E.L., Iwanaga, R., Jedlicka, P., Abbey, N. S., Chodosh, L. A., Heichman, K. A., Welm, A. L. & Ford, H. L. Six1 expands the mouse mammary epithelial stem/progenitor cell pool and induces mammary tumors that undergo epithelial-mesenchymal transition. *J Clin Invest* 119, 2663-2677 (2009).
- Micalizzi, D.S., Christensen, K. L., Jedlicka, P., Coletta, R. D., Baron, A. E., Harrell, J. C., Horwitz, K. B., Billheimer, D., Heichman, K. A., Welm, A. L., Schiemann, W. P. & Ford, H. L. The Six1 homeoprotein induces human mammary carcinoma cells to undergo epithelial-mesenchymal transition and metastasis in mice through increasing TGF-beta signaling. *Journal of Clinical Investigation* 119, 2678-2690 (2009).
- Mummler, C., Burgy, O., Hermann, S., Mutze, K., Gunther, A., & Konigshoff, M. Cell-specific expression of runt-related transcription factor 2 contributes to pulmonary fibrosis. *Faseb Journal* 32, 703-716 (2018).

- Nakamura, Y., Esnault, S., Maeda, T., Kelly, E. A. B., Malter, J. S. & Jarjour, N. N. Ets-1 regulates TNF-alpha-induced matrix metalloproteinase-9 and tenascin expression in primary bronchial fibroblasts. *Journal of Immunology* 172, 1945-1952 (2004).
- Nakayama, T., Ito, M., Ohtsuru, A., Naito, S., Nakashima, M., Fagin, J. A., Yamashita, S. & Sekine, I. Expression of the Ets-1 proto-oncogene in human gastric carcinoma - Correlation with tumor invasion. *American Journal of Pathology* 149, 1931-1939 (1996).
- Ocana, O.H., Corcoles, R., Fabra, A., Moreno-Bueno, G., Acloque, H., Vega, S., Barrallo-Gimeno, A., Cano, A. & Nieto, M. A. Metastatic Colonization Requires the Repression of the Epithelial-Mesenchymal Transition Inducer Prrx1. *Cancer Cell* 22, 709-724 (2012).
- Overholtzer, M., Zhang, J., Smolen, G. A., Muir, B., Li, W., Sgroi, D. C., Deng, C. X., Brugge, J. S. & Haber, D. A. Transforming properties of YAP, a candidate oncogene on the chromosome 11q22 amplicon. *Proceedings of the National Academy of Sciences of the United States of America* 103, 12405-12410 (2006).
- Perez-Moreno, M.A., Locascio, A., Rodrigo, I., Dhondt, G., Portillo, F., Nieto, M. A. & Cano, A. A new role for E12/E47 in the repression of E-cadherin expression and epithelial-mesenchymal transitions. *Journal of Biological Chemistry* 276, 27424-27431 (2001).
- Petrovics, G., Liu, A. J., Shaheduzzaman, S., Furasato, B., Sun, C., Chen, Y. M., Nau, M., Ravindranath, L., Chen, Y. D., Dobi, A., Srikantan, V., Sesterhenn, I. A., McLeod, D. G., Vahey, M., Moul, J. W. & Srivastava, S. Frequent overexpression of ETS-related gene-1 (ERG1) in prostate cancer transcriptome. *Oncogene* 24, 3847-3852 (2005).
- Pratap, J., Javed, A., Languino, L. R., van Wijnen, A. J., Stein, J. L., Stein, G. S. & Lian, J. B. The Runx2 Osteogenic Transcription Factor Regulates Matrix Metalloproteinase 9 in Bone Metastatic Cancer Cells and Controls Cell Invasion. *Molecular and Cellular Biology* 25, 8581-8591 (2005).
- Reichmann, E., Schwarz, H., Deiner, E. M., Leitner, I., Eilers, M., Berger, J., Busslinger, M. & Beug, H. Activation of an inducible c-FosER fusion protein causes loss of epithelial polarity and triggers epithelial-fibroblastoid cell conversion. *Cell* 71, 1103-1116 (1992).
- Rodriguez, M., Aladowicz, E., Lanfrancone, L. & Goding, C.R. Tbx3 represses E-cadherin expression and enhances melanoma invasiveness. *Cancer Research* 68, 7872-7881 (2008).
- Saunders, L.R. & McClay, D.R. Sub-circuits of a gene regulatory network control a developmental epithelial-mesenchymal transition. *Development* 141, 1503-1513 (2014).
- Seo, E., Kim, W. Y., Hur, J., Kim, H., Nam, S. A., Choi, A., Kim, Y. M., Park, S. H., Chung, C., Kim, J., Min, S., Myung, S. J., Lim, D. S. & Kim, Y. K. The Hippo-Salvador signaling pathway regulates renal tubulointerstitial fibrosis. *Scientific Reports* 6 (2016).
- Singh, R. et al. Tbx2 and Tbx3 induce atrioventricular myocardial development and endocardial cushion formation. *Cell Mol Life Sci* 69, 1377-1389 (2012).
- Sipos, F. & Galamb, O. Epithelial-to-mesenchymal and mesenchymal-to-epithelial transitions in the colon. *World J Gastroentero* 18, 601-608 (2012).

- Slattery, C., McMorro, T. & Ryan, M.P. Overexpression of E2A proteins induces epithelial-mesenchymal transition in human renal proximal tubular epithelial cells suggesting a potential role in renal fibrosis. *Febs Letters* 580, 4021-4030 (2006).
- Smith, J.C., Boone, B.E., Opalenik, S.R., Williams, S.M. & Russell, S.B. Gene profiling of keloid fibroblasts shows altered expression in multiple fibrosis-associated pathways. *Journal of Investigative Dermatology* 128, 1298-1310 (2008).
- Sobrado, V.R., Moreno-Bueno, G., Cubillo, E., Holt, L. J., Nieto, M. A., Portillo, F. & Cano, A. The class I bHLH factors E2-2A and E2-2B regulate EMT. *Journal of Cell Science* 122, 1014-1024 (2009).
- Sun, S., Sun, W., Xia, L., Liu, L., Du, R., He, L., Li, R., Wang, H & Huang, C. The T-box transcription factor Brachyury promotes renal interstitial fibrosis by repressing E-cadherin expression. *Cell Commun Signal* 12, 76 (2014).
- Sunami, Y., Leithauser, F., Gul, S., Fiedler, K., Guldiken, N., Espenlaub, S., Holzmann, K. H., Hipp, N., Sindrilari, A., Luedde, T., Baumann, B., Wissel, S., Kreppel, F., Schneider, M., Scharffetter-Kochanek, K., Kochanek, S., Strnad, P. & Wirth, T. Hepatic activation of IKK/NFkappaB signaling induces liver fibrosis via macrophage-mediated chronic inflammation. *Hepatology* 56, 1117-1128 (2012).
- Tavares, A.L.P., Brown, J.A., Ulrich, E.C., Dvorak, K. & Runyan, R.B. Runx2-I is an Early Regulator of Epithelial-Mesenchymal Cell Transition in the Chick Embryo. *Dev Dyn* (2017).
- Tiwari, N., Meyer-Schaller, N., Arnold, P., Antoniadis, H., Pachkov, M., van Nimwegen, E. & Christofori, G. Klf4 Is a Transcriptional Regulator of Genes Critical for EMT, Including Jnk1 (Mapk8). *Plos One* 8 (2013).
- Tobin, N.P., Sims, A.H., Lundgren, K.L., Lehn, S. & Landberg, G. Cyclin D1, Id1 and EMT in breast cancer. *Bmc Cancer* 11, 417 (2011).
- Venkov, C.D., Link, A. J., Jennings, J. L., Plieth, D., Inoue, T., Nagai, K., Xu, C., Dimitrova, Y. N., Rauscher, F. J. & Neilson, E. G. A proximal activator of transcription in epithelial-mesenchymal transition. *Journal of Clinical Investigation* 117, 482-491 (2007).
- Wang, X., Zheng, M., Liu, G., Xia, W., McKeown-Longo, P. J., Hung, M. C. & Zhao, J. Kruppel-like factor 8 induces epithelial to mesenchymal transition and epithelial cell invasion. *Cancer Res* 67, 7184-7193 (2007).
- Wensing, L.A. & Campos, A.H. TBX3, a downstream target of TGF-beta 1, inhibits mesangial cell apoptosis. *Experimental Cell Research* 328, 340-350 (2014).
- Wu, X., Chen, H., Parker, B., Rubin, E., Zhu, T., Lee, J. S., Argani, P. & Sukumar, S. HOXB7, a homeodomain protein, is overexpressed in breast cancer and confers epithelial-mesenchymal transition. *Cancer Res* 66, 9527-9534 (2006).
- Xiang, G., Yi, Y., Weiwei, H. & Weiming, W. TGIF1 promoted the growth and migration of cancer cells in nonsmall cell lung cancer. *Tumour Biol* 36, 9303-9310 (2015).
- Xiao, L., Zhou, X., Liu, F. Y., Hu, C., Zhu, X. J., Luo, Y., Wang, M., Xu, X. X., Yang, S. K., Kanwar, Y. S. & Sun, L. MicroRNA-129-5p modulates epithelial-to-mesenchymal transition by targeting SIP1 and SOX4 during peritoneal dialysis. *Laboratory Investigation* 95, 817-832 (2015).

- Yuan, H., Kajiyama, H., Ito, S., Yoshikawa, N., Hyodo, T., Asano, E., Hasegawa, H., Maeda, M., Shibata, K., Hamaguchi, M., Kikkawa, F. & Senga, T. ALX1 induces snail expression to promote epithelial-to-mesenchymal transition and invasion of ovarian cancer cells. *Cancer Res* 73, 1581-1590 (2013).
- Zhang, J., Liang, Q., Lei, Y., Yao, M., Li, L., Gao, X., Feng, J., Zhang, Y., Gao, H., Liu, D. X., Lu, J. & Huang, B. SOX4 induces epithelial-mesenchymal transition and contributes to breast cancer progression. *Cancer Res* 72, 4597-4608 (2012).