



ERJ European Respiratory Journal Eur Respir J 0903-1936/1399-3003 European Respiratory Society 10.1183/13993003.01708-2020 ERJ-01708-2020 AGORA RESEARCH LETTER

Clinical characteristics of COVID-19 and active tuberculosis co-infection in an Italian reference hospital. C. STOCHINO ET AL. RESEARCH LETTER Clinical characteristics of COVID-19 and active TB co-infection

Stochino Claudia¹, Villa Simone², Zucchi Patrizia³, Parravicini Pierpaolo³, Gori Andrea^{2,4}, Raviglione Mario Carlo²,

¹Phthysiology Unit, Sondalo Hospital, ASST Valtellina e Alto Lario, Sondrio, Italy. ²Centre for Multidisciplinary Research in Health Science, University of Milan, Milan, Italy. ³Dept of Internal Medicine, ASST Valtellina e Alto Lario, Sondrio, Italy. ⁴Infectious Diseases Unit, Dept of Internal Medicine, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy.

Correspondence: Simone Villa, Centre for Multidisciplinary Research in Health Science, University of Milan, Via Francesco Sforza 35, 20122, Milan, Italy. E-mail: simone.villa@unimi.it
2020202056010052020250520202020

To the Editor:

Coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), emerged in Wuhan, China, in December 2019. In February 2020, an outbreak was detected in the Lombardy region of Italy, resulting in the first major outbreak outside Asia [1].

