

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Road Traffic Noise Exposure and Filled Prescriptions for Antihypertensive Medication: A Danish Cohort Study

Jesse D. Thacher, Aslak H. Poulsen, Nina Roswall, Ulla Hvidtfeldt, Ole Raaschou-Nielsen, Steen Solvang Jensen, Matthias Ketzel, Jørgen Brandt, Kim Overvad, Anne Tjønneland, Thomas Münzel, and Mette Sørensen

Table of Contents

Figure S1. Graphical evaluation of estimated exposure-response curves (solid lines) with 95% confidence intervals (dashed lines) for exposures and selected covariates.

Figure S2. Study Population.

Table S1. Correlation matrix of road traffic noise and air pollution.

Table S2. Associations between residential exposure to traffic noise (per 10 dB) and filled prescriptions for antihypertensive medication (excluding diuretics).

Table S3. Associations between residential exposure to traffic noise (per 10 dB) and filled prescriptions for antihypertensive medication adjusted for air pollution.

Table S4. Association between categories of exposure to road traffic (L_{den}) at the most exposed façade and filled prescription for antihypertensive medication.

Table S5. Association between categories of exposure to road traffic (L_{den}) at the least exposed façade and filled prescriptions for antihypertensive medication.

Figure S3. Distribution of residential exposure to road traffic (L_{den}) at baseline.