Supplementary table 1 Study information

| Country | Study location | Stratification for sampling ^a | Recruitment | Dates | Age range (years) | Total sample ^e (n) | Compliance | Deprivation measure | Reference |
|---------|--|---|--|----------------|-------------------------|-------------------------------|------------|------------------------|-----------|
| England | Blackburn with Darwen (North West England) | Blackburn with Darwen Local Authority ^b | Study information letter sent to randomly | 2012 (Aug-Sep) | 18-70 ^d | 1500 | 70.4% | IMD 2010 | 1 |
| England | National sample | English Administrative Regions (n=10) ^c | selected households; households not opting out upon receipt of | 2013 (Apr-Jul) | 18-69 | 4010 | 53.5% | IMD 2010 | 2 |
| England | Luton, Hertfordshire Northamptonshire (South East England) | Luton, Hertfordshire Northamptonshire (n=3) | letter visited by researchers | 2015 (Jun-Sep) | 18-69 | 5623 | 55.8% | IMD 2011 | 3 |
| Wales | National sample | Welsh Health Regions (n=7) | | 2017 (Mar-Jun) | 18-69 | 2497 | 58.5% | WIMD 2014 | 4 |
| Wales | National sample | Welsh Health Regions (n=7) | Households in sampled areas randomly selected by researchers | 2015 (Feb-May) | 18-69 | 2028 | 49.1% | WIMD 2014 | 5 |

^aLower Super Output Area level stratification by deprivation quintile. ^bNo sub-regional stratification was undertaken in Blackburn with Darwen due to the relatively small size of the sample area. ^cLondon was split into Inner and Outer London for regional sampling. ^dIndividuals aged 70 years were excluded from the sample for consistency. ^eIndividuals not completing all questions on variables of interest were excluded. IMD, Index of Multiple Deprivation; WIMD, Welsh Index of Multiple Deprivation.

- 1. Bellis MA, Lowey H, Leckenby N, Hughes K, Harrison D. Adverse childhood experiences: retrospective study to determine their impact on adult health behaviours and health outcomes in a UK population. J Public Health 2014;36:81-91.
- 2. Bellis MA, Hughes K, Leckenby N, Hardcastle K, Perkins C, Lowey H. Measuring mortality and the burden of adult disease associated with Adverse Childhood Experiences in England: a national survey. J Public Health 2015;37:445-54.
- 3. Ford K, Butler N, Hughes K, Quigg Z, Bellis MA. Adverse childhood experiences in Hertfordshire, Luton and Northamptonshire. Liverpool: Centre for Public Health, Liverpool John Moores University, 2016. https://www.researchgate.net/publication/302589403 Adverse Childhood Experiences ACEs in Hertfordshire Luton and Northamptonshire (accessed 16 October 2019).
- 4 Hughes K, Ford K, Davies AR, Homolova L, Bellis MA. Sources of resilience and their moderating relationships with harms from adverse childhood experiences. Wrexham: Public Health Wales, 2018.

 <a href="https://www.researchgate.net/publication/322697404_Sources_of_resilience_and_their_moderating_relationships_with_harms_from_adverse_childhood_experiences_Report_1_M ental Illness (accessed 16 October 2019).
- 5 Bellis MA, Ashton K, Hughes K, Ford K, Bishop J, Paranjothy S. Adverse childhood experiences and their impact on health-harming behaviours in the Welsh population. Cardiff: Public Health Wales, 2016. https://www.researchgate.net/publication/290433553 Adverse Childhood Experiences and their impact on health-harming behaviours in the Welsh adult population (accessed 16 October 2019).

1

Supplementary Table 2 ACE and health outcome questions used

| ACE questions. All A | ACE questions were preceded by the statement "While you v | were growing up, before the age of 18" |
|----------------------|---|---|
| ACE | Question | Qualifying responses |
| Physical abuse | How often did a parent or adult in your home ever hit, beat, kick, or physically hurt you in any way? This does not include gentle smacking for punishment? | Once; more than once |
| Verbal abuse | How often did a parent or adult in your home ever swear at you, insult you, or put you down? | More than once |
| Sexual abuse | How often did anyone at least 5 years older than you (including adults) evertouch you sexually?try to make you touch them sexually?force you to have any type of sexual intercourse (oral, anal, or vaginal)? | Once or more than once to any of the questions |
| Parental separation | Were your parents ever separated or divorced? | Yes |
| Domestic violence | How often did your parents or adults in your home ever slap, hit, kick, punch, or beat each other up? | Once; more than once |
| Mental illness | Did you live with anyone who was depressed, mentally ill, or suicidal? | Yes |
| Alcohol abuse | Did you live with anyone who was a problem drinker or alcoholic? | Yes |
| Drug abuse | Did you live with anyone who used illegal street drugs or who abused prescription medications? | Yes |
| Incarceration | Did you live with anyone who served time or was sentenced to serve time in a prison or young offenders' institution? | Yes |
| Health outcome | Question | Qualifying responses |
| Alcohol use | How often do you have a drink containing alcohol? How many standard drinks containing alcohol do you have on a typical day when you drink? - responses used to estimate daily consumption | Consumption of 12g or more alcohol per day |
| Smoking | In terms of smoking tobacco, which of the following best describes you? | I smoke daily; I smoke occasionally but not daily |
| Drug use | How often, if ever, have you used heroin or crack cocaine? | Used but not in the last 12 months; Used in the past 12 months |
| High BMI | What is your height? (in feet/inches or metres/centimetres) What is your weight? (in stone/pound, kilograms or pounds) - Responses used to calculate BMI | BMI 25.0 or higher |
| Depression | Are you currently or have you ever been treated for depression? | Yes, currently; Yes, in the past |
| Anxiety | Are you currently or have you ever been treated for anxiety? | Yes, currently; Yes, in the past |
| Other mental illness | Are you currently or have you ever been treated for another mental illness? | No, never; Yes, currently; Yes, in the past |
| Victim of violence | How many times have you been physically hit in the past 12 months? Or (in Wales, 2017): In the past 12 months, have you been physically hit by someone else? | Once; 2 or 3 times; More than 3 times Yes |
| Cancer | Has a doctor or nurse ever told you that you have Cancer? | Yes |
| Type 2 diabetes | Has a doctor or nurse ever told you that you have Type 2 diabetes? | Yes |
| Heart disease | Has a doctor or nurse ever told you that you have Coronary heart disease or heart attack? | Yes |
| Stroke | Has a doctor or nurse ever told you that you have Stroke? | Yes |
| Respiratory disease | Has a doctor or nurse ever told you that you have Respiratory disease such as Chronic bronchitis/ Emphysema/ Chronic Obstructive Pulmonary Disease? | Yes |

ACE, adverse childhood experience; BMI, body mass index.

Supplementary Table 3 Outcomes measured across studies

| Supplementary Table 5 | | | England | England | Wales | Wales |
|-----------------------|---------|------------|------------|---------|------------|------------|
| | | England | (South | (North | 2015 | 2017 |
| | All | (national) | East) | West) | (national) | (national) |
| Total sample (n) | 15285 | 3885 | 5454 | 1421 | 2028 | 2497 |
| Alcohol use | n=12736 | | | | | |
| Missing (n) | 52 | 18 | 7 | 23 | 4 | - |
| Yes (%) | 13.6 | 16.3 | 10.0 | 15.3 | 17.1 | - |
| Smoking | n=15281 | | | | | |
| Missing (n) | 4 | 0 | 0 | 4 | 0 | 0 |
| Yes (%) | 26.3 | 26.9 | 23.1 | 37.1 | 27.9 | 25.3 |
| Drug use | n=15246 | | | | | |
| Missing (n) | 39 | 3 | 19 | 14 | 2 | 1 |
| Yes (%) | 2.4 | 2.2 | 2.0 | 3.1 | 4.4 | 1.6 |
| High BMI | n=11527 | | | | | |
| Missing (n) | 1261 | 424 | 481 | 16 | 340 | - |
| Yes (%) | 49.8 | 50.9 | 49.0 | 45.7 | 53.2 | - |
| Depression | n=2496 | | | | | |
| Missing (n) | 1 | - | - | - | - | 1 |
| Yes (%) | 29.2 | = | - | - | - | 29.2 |
| Anxiety | n=2493 | | | | | |
| Missing (n) | 4 | - | - | - | - | 4 |
| Yes (%) | 25.0 | - | - | - | - | 25.0 |
| Other mental illness | n=2491 | | | | | |
| Missing (n) | 6 | - | - | - | - | 6 |
| Yes (%) | 6.0 | - | - | - | - | 6.0 |
| Victim of violence | n=15267 | | | | | |
| Missing (n) | 18 | 2 | 10 | 2 | 4 | 0 |
| Yes (%) | 5.1 | 5.3 | 3.7 | 6.3 | 9.1 | 4.1 |
| Cancer | n=12765 | | | _ | _ | |
| Missing (n) | 23 | 4 | 12 | 7 | 0 | - |
| Yes (%) | 3.7 | 4.4 | 2.9 | 3.2 | 4.8 | - |
| Type 2 Diabetes | n=12769 | | | | | |
| Missing (n) | 19 | 3 | 12 | 4 | 0 | - |
| Yes (%) | 5.1 | 4.8 | 4.7 | 7.3 | 5.3 | - |
| Heart disease | n=12773 | | | | | |
| Missing (n) | 15 | 1 | 12 | 2 | 0 | - |
| Yes (%) | 2.7 | 3.2 | 2.3 | 4.2 | 1.5 | - |
| Stroke | n=12773 | _ | . - | _ | _ | |
| Missing (n) | 19 | 1 | 13 | 5 | 0 | - |
| Yes (%) | 1.1 | 1.1 | 1.0 | 1.8 | 0.8 | - |
| Respiratory disease | n=12766 | _ | | _ | _ | |
| Missing (n) | 22 | 0 | 15 | 7 | 0 | - |
| Yes (%) | 2.8 | 3.5 | 1.9 | 5.2 | 2.1 | - |

⁻ outcome not measured in survey. BMI, body mass index

Supplementary Table 4: Study outcome and matched Global Burden of Disease (GBD) category

| Outcome | GBD category matched (ID) | | | | | |
|----------------------|---|--|--|--|--|--|
| Risk factors | | | | | | |
| Alcohol use | Alcohol use (102) | | | | | |
| Smoking | Smoking (99) | | | | | |
| Drug use | Drug use (103) | | | | | |
| High BMI | High body-mass index (108) | | | | | |
| Causes of ill health | | | | | | |
| Depression | Major depressive disorder (586) | | | | | |
| Anxiety | Anxiety disorders (571) | | | | | |
| Other mental illness | Mental disorders (558), excluding major depressive disorder (586) and anxiety | | | | | |
| | disorders (571) | | | | | |
| Violence | Interpersonal violence (724) | | | | | |
| Cancer | Neoplasms (410) | | | | | |
| Type 2 diabetes | Diabetes mellitus type 2 (976) | | | | | |
| Heart disease | Ischaemic heart disease (493), hypertensive heart disease (498) | | | | | |
| Stroke | Stroke (494) | | | | | |
| Respiratory disease | Chronic respiratory diseases (508), excluding asthma (515) | | | | | |

ID, identification; BMI, body mass index

Supplementary Table 5 Sample demographics and ACE count prevalence

| | • | England | England | England | Wales 2015 | Wales 2017 |
|--------------------------|-------|------------|--------------|--------------|------------|-------------------|
| | All | (national) | (South East) | (North West) | (national) | (national) |
| Total sample (n) | 15285 | 3885 | 5454 | 1421 | 2028 | 2497 |
| Gender (%) | | | | | | |
| Male | 45.1 | 45.0 | 44.7 | 39.9 | 49.8 | 45.3 |
| Female | 54.9 | 55.0 | 55.3 | 60.1 | 50.2 | 54.7 |
| Age group (%) | | | | | | |
| 18-29 | 21.9 | 21.0 | 20.6 | 24.6 | 30.4 | 17.9 |
| 30-39 | 19.9 | 19.9 | 22.5 | 21.5 | 14.2 | 18.4 |
| 40-49 | 20.3 | 20.5 | 20.6 | 22.2 | 17.8 | 20.1 |
| 50-59 | 17.7 | 18.0 | 17.0 | 14.6 | 17.5 | 20.6 |
| 60-69 | 20.2 | 20.7 | 19.3 | 17.0 | 20.2 | 23.1 |
| Ethnicity (%) | | | | | | |
| White | 85.7 | 86.3 | 80.6 | 71.3 | 95.4 | 96.4 |
| Other | 14.3 | 13.7 | 19.4 | 28.7 | 4.6 | 3.6 |
| Deprivation quintile (%) | | | | | | |
| (least deprived) 1 | 21.7 | 20.1 | 28.5 | 5.5 | 21.7 | 18.7 |
| 2 | 19.1 | 19.5 | 20.2 | 10.2 | 19.4 | 20.9 |
| 3 | 19.9 | 19.7 | 20.8 | 8.1 | 19.4 | 25.1 |
| 4 | 19.2 | 19.9 | 20.0 | 14.5 | 18.7 | 19.3 |
| (most deprived) 5 | 20.1 | 20.7 | 10.5 | 61.7 | 20.7 | 15.9 |
| ACE count (%) | | | | | | |
| 0 | 55.3 | 53.6 | 58.4 | 54.4 | 54.4 | 52.1 |
| 1 | 19.7 | 22.7 | 18.2 | 20.0 | 19.0 | 18.9 |
| 2-3 | 15.3 | 15.4 | 15.3 | 15.8 | 13.0 | 17.1 |
| ≥4 | 9.6 | 8.3 | 8.1 | 9.8 | 13.6 | 11.9 |

ACE, Adverse childhood experience.

Supplementary Table 6 Unadjusted proportion reporting each outcome by ACE count category

| | | ACE count | | | | |
|----------------------|-------|-----------|-------|----------|---------|--|
| | n | 0 ACEs | 1 ACE | 2-3 ACEs | ≥4 ACEs | |
| Alcohol use | 12736 | 11.3 | 14.6 | 15.9 | 22.0 | |
| Smoking | 15281 | 20.3 | 26.3 | 32.9 | 50.6 | |
| Drug use | 15246 | 0.9 | 1.8 | 3.4 | 10.5 | |
| High BMI | 11527 | 49.3 | 49.8 | 51.1 | 50.6 | |
| Depression | 2496 | 18.4 | 29.4 | 41.3 | 59.1 | |
| Anxiety | 2493 | 16.0 | 24.2 | 36.5 | 49.8 | |
| Other mental illness | 2491 | 2.8 | 4.9 | 9.9 | 16.4 | |
| Violence | 15267 | 2.5 | 3.8 | 8.3 | 18.2 | |
| Cancer | 12765 | 3.5 | 3.5 | 3.7 | 5.1 | |
| Type 2 diabetes | 12769 | 5.0 | 5.0 | 5.3 | 5.6 | |
| Heart disease | 12773 | 2.5 | 2.5 | 3.2 | 3.1 | |
| Stroke | 12769 | 0.9 | 1.1 | 1.3 | 1.8 | |
| Respiratory disease | 12766 | 2.2 | 2.7 | 4.0 | 4.7 | |

ACE, adverse childhood experience; BMI, body mass index