

**Supplemental Table 1: Showing the relationship between location, response time and percentage survival to hospital admission for those patients who received a shock. ‘Public’ included public streets, parks, hotels, restaurants and recreational facilities excluding those locations reported otherwise. Medical facilities were non-hospital treatment centres, such as general practitioner offices, dental offices or residential care facilities, where the PAD is available for use by lay-rescuers.**

Location	Events (n)	Mean (SD) Patient Age	Events with Known Response Times (n)	Median (IQR) Response Time (minutes)	Events with response time 5-minutes or less (N and %)	Events with Known Survival to Hospital Admission (n)	Survived to Hospital Admission n (%)
Public	141	60 (13)	82	3 (IQR 2 to 5)	56 (68.3%)	134	76 (56.7%)
Sports Facility	76	53 (16)	38	2 (IQR 1 to 5)	28 (73.7%)	75	68 (90.7%)
Home	64	62 (13)	38	5 (IQR 4 to 6)	14 (36.8%)	43	18 (41.9%)
Office	30	55 (8)	16	4 (IQR 2 to 9)	8 (50.0%)	28	20 (71.4%)
Medical Facility	21	64 (13)	9	3 (IQR 2 to 5)	5 (55.6%)	19	11 (57.9%)
School/University	7	21 (19)	6	5 (IQR 2 to 5)	2 (33.3%)	7	6 (85.7%)
Unknown	15	70 (12)	1	3 (IQR 3 to 3)	1 (100.0%)	6	4 (66.7%)
Total	354	58 (15)	190	4 (IQR 2 to 5)	114 (60.0%)	312	203 (65.1%)