

Table S2: Risk of Ischaemic Heart Disease and Myocardial Infarction by Exposure Status Adjusted for Number of GP Consultations

	IHD			MI				
	Male		Female		Male		Female	
	HR (95%CI)		HR (95%CI)		HR (95%CI)		HR (95%CI)	
ILD	1.95 (1.64-2.32)		1.58 (1.19-2.08)		1.87 (1.57-2.22)		1.72 (1.34-2.22)	
PS	1.40 (0.97-2.01)		0.97 (0.61-1.55)		2.00 (1.45-2.75)		0.95 (0.59-1.51)	
PF	2.06 (1.71-2.48)		1.97 (1.43-2.72)		1.75 (1.44-2.12)		2.16 (1.63-2.86)	
Age								
	IHD			MI				
	ILD	PS	PF	ILD	PS	PF		
<50	1.73 (0.98-3.06)	1.57 (0.87-2.84)	2.10 (0.61-7.29)	1.84 (1.11-3.07)	1.94 (1.15-3.28)	0.92 (0.21-4.06)		
50-59	1.42 (0.90-2.28)	0.91 (0.50-1.67)	2.39 (1.31-4.36)	1.96 (1.25-3.06)	1.90 (1.12-3.24)	1.64 (0.86-3.10)		
60-69	2.17 (1.65-2.86)	0.96 (0.56-1.67)	2.74 (2.04-3.68)	2.10 (1.58-2.80)	0.69 (0.35-1.35)	2.80 (2.07-3.78)		
70-79	1.94 (1.53-2.46)	0.86 (0.38-1.94)	2.10 (1.64-2.68)	2.03 (1.61-2.57)	1.46 (0.78-2.75)	2.08 (1.63-2.66)		
>80	1.30 (0.87-1.95)	0.82 (0.10-6.69)	1.33 (0.88-1.99)	1.09 (0.77-1.55)	1.18 (0.33-4.28)	1.09 (0.76-1.56)		
All analyses adjusted for CKD, HTN, DM, HLD, BMI, exposure to smoking and alcohol, IMD, family history of cardiovascular disease and exposure to anti-hypertensive, anti-platelet, lipid-lowering drugs and no of GP consultations								
CI=Confidence Interval; CKD=Chronic Kidney Disease; DM=Diabetes Mellitus; HLD=Hyperlipidaemia; HR= adjusted Hazard Ratio; HTN=Hypertension; IHD=Ischaemic Heart Disease; ILD=Inflammatory Lung Disease; IMD=Index of Multiple Deprivation; MI=Myocardial Infarction; PF=Pulmonary Fibrosis; PS=Pulmonary Sarcoidosis								