

| Characteristics | Non-DM | DM | SMD |
|--------------------------------------|-------------------|------------------|------------|
| | (n=1309.1) | (n=602.5) | |
| Age (years) | 64.02 (11.14) | 64.16 (10.08) | 0.013 |
| Men | 0.70 (0.46) | 0.70 (0.46) | 0.002 |
| Body-mass index (kg/m ²) | 24.41 (3.15) | 24.41 (3.23) | 0.001 |
| Hypertension | 0.55 (0.50) | 0.55 (0.50) | 0.01 |
| Hyperlipidemia | 0.20 (0.40) | 0.20 (0.40) | 0.012 |
| Current smoker | 0.35 (0.48) | 0.35 (0.48) | 0.017 |
| Atrial fibrillation | 0.02 (0.15) | 0.02 (0.14) | 0.014 |
| Previous MI | 0.02 (0.14) | 0.02 (0.15) | 0.022 |
| Previous PCI | 0.09 (0.28) | 0.09 (0.29) | 0.004 |
| Previous CABG | 0.01 (0.11) | 0.01 (0.11) | 0.009 |
| Renal failure | 0.03 (0.18) | 0.03 (0.18) | 0.002 |
| Cerebrovascular disease | 0.08 (0.27) | 0.08 (0.27) | 0.001 |
| Ejection fraction (%) | 56.8 (10.7) | 56.94 (10.5) | 0.011 |
| Clinical presentation | | | 0.001 |
| Stable angina | 0.31 (0.46) | 0.31 (0.46) | |
| Unstable angina | 0.29 (0.45) | 0.29 (0.46) | |
| NSTEMI | 0.20 (0.40) | 0.20 (0.40) | |
| STEMI | 0.20 (0.40) | 0.20 (0.40) | |
| Discharge medications | | | |
| Aspirin | 0.99 (0.11) | 0.99 (0.12) | 0.007 |
| ADP receptor antagonist | 0.93 (0.26) | 0.93 (0.25) | 0.018 |

| | | | |
|-------------------------|-------------|-------------|-------|
| Cilostazol | 0.19 (0.39) | 0.19 (0.39) | 0.002 |
| β-blocker | 0.75 (0.43) | 0.74 (0.44) | 0.01 |
| Calcium channel blocker | 0.18 (0.39) | 0.19 (0.39) | 0.008 |
| ACE inhibitor or ARB | 0.66 (0.47) | 0.65 (0.48) | 0.01 |
| Statin | 0.81 (0.39) | 0.81 (0.40) | 0.009 |

Data are shown as mean (SD) for continuous variables and absolute numbers (percentage) for dichotomous variables. Abbreviations: CABG, coronary artery bypass grafting; CAD, coronary artery disease; CHF, congestive heart failure; NSTEMI, non-ST-elevation myocardial infarction; MI, myocardial infarction; PCI, percutaneous coronary intervention; STEMI, ST-elevation MI; SMD, standardization mean differences