

Table S1. Primers used for qRT-PCR analysis of mRNA expression and immunoprecipitated chromatin.

qRT-PCR primers for gene expression analysis		
Gene	Forward(5'→3')	Reverse (5'→3')
<i>PFKFB3</i>	ATTGCGGTTTCGATGCCAC	GCCACAACGTAGGGTCGT
<i>PFKP</i>	CGCCTACCTCAACGTGGTG	ACCTCCAGAACGAAGGTCC
<i>HK2</i>	TGCCACCAGACTAAACTAGACG	CCCGTGCCCACAATGAGAC
<i>PKM1</i>	CGAGCCTCAAGTCACTCCA	GTGAGCAGACCTGCCAGAC
<i>PKM2</i>	ATAACGCCTACATGGAAAAGTGT	TAAGCCCATCATCCACGTAGA
18s	GATGGGCGGCGGAAAATAG	GCGTGGATTCTGCATAATGGT
qRT-PCR primers for analysis of immunoprecipitated chromatin		
<i>TNF</i> promoter R1	CAGGCAGGTTCTCTTCCCT	GCTTCAGTGCTCATGGTGT
<i>TNF</i> promoter R2	AGAGGACCAGCTAACAGAGGGA	AGCTTGTCAAGGGATGTGG
<i>TNF</i> promoter R3	GTGCTTGTTCCTCAGCCTCT	ATCACTCAAAGTGCAGCAG
<i>TNF</i> promoter R4	TGTCTGGCACACAGAACAGA	CCCTGAGGTGTCTGGTTTC
<i>TNF</i> promoter R5	TGATGGTAGGCAGAACATTGG	ACTAAGGCCTGTGCTGTTCC
<i>IL6</i> promoter R1	TCGTGCATGACTTCAGCTT	GCGCTAACAGAACAGAACAC
<i>IL6</i> promoter R2	AGGGAGAGCCAGAACACAGA	GAGTTCCCTGACTCCATCG
<i>PFKFB3</i> promoter R1	TCTTCTACTCGGGCGATAA	GGAGCTGGACTGAAGTGGAC
<i>PFKFB3</i> promoter R2	TCAAGGTTCCCAGTCTTGG	CACCTCGTCCTGTCCACTT
<i>PFKFB3</i> promoter R3	ACTGCGTCTCTGCTCTCCTC	GCTGCTGTTATCCCCTCGT
<i>HK2</i> promoter R1	GAGCTCAATTCTGTGTGGAGT	ACTTCTTGAGAACTATGTACCCCTT
<i>PFKP</i> promoter R1	CGAAGGCGATGGGGTGAC	CATCGCTTCGCCACCTTTC