

1 **SUPPLEMENTARY INFORMATION**

2

3 **Table S1.** Summary of screen failure reasons for ineligible patients (n=26)

4

Inclusion/exclusion criteria	Screen failure reason	Screen failures n (%) ^a
Patients must have active axSpA disease at screening as defined by BASDAI \geq 4 and spinal pain \geq 4 (from BASDAI item 2).	Ineligibility	10 (38)
Patients must have a documented history of AAU diagnosed by an ophthalmologist and have at least 2 AAU flares in the past, of which at least 1 AAU flare was in the last 12 months prior to baseline.	Ineligibility	5 (19)
Patients must be HLA-B27 positive.	Ineligibility	3 (12)
Patients must not have any other inflammatory arthritis.	Ineligibility	1 (4)
Patient has a condition that may interfere with the AAU assessment.	Ineligibility	1 (4)
Patient has received intraocular or periocular corticosteroids within 90 days prior to the baseline visit.	Ineligibility	3 (12)
Patients must not have used any forbidden co-medication as per protocol.	Ineligibility	1 (4)
Patients have a history of chronic or recurrent infections, excluding uveitis.	Adverse event	1 (4)
Patients have current acute or chronic viral hepatitis B or C, or human immunodeficiency virus infection.	Ineligibility	1 (4)
Patients with current or history of active infection with histoplasma, coccidioides, para coccidioides, pneumocystis, nontuberculous mycobacteria, blastomyces, or aspergillus.	Ineligibility	1 (4)
Patients with significant laboratory abnormalities.	Ineligibility	1 (4)

5 [a] The denominator is the total number of screen failures (n=26). Patients could have more than one reason for screen failure. AAU: acute anterior uveitis;
 6 axSpA: axial spondyloarthritis; BASDAI: Bath Ankylosing Spondylitis Disease Activity Index.

7