

Supplementary information

Non-canonical chemical feedback self-limits nitric oxide-cyclic GMP signaling in health and disease

Vu Thao-Vi Dao^{1,2}, Mahmoud H. Elbatreek^{1,3*}, Martin Deile⁴, Pavel I. Nedvetsky⁵, Andreas Güldner⁶, César Ibarra-Alvarado⁷, Axel Gödecke⁸ and Harald H.H.W. Schmidt^{1*}

¹Department of Pharmacology and Personalised Medicine, MeHNS, FHML, Maastricht University, Maastricht, The Netherlands;

²Department of Pharmacology, Johannes Gutenberg University, Mainz, Germany

³Department for Pharmacology and Toxicology, Faculty of Pharmacy, Zagazig University, Zagazig, Egypt;

⁴Asklepios-Klinik Radeberg, Dresden, Germany;

⁵Universitätsklinikum Münster, Medical Cell Biology, Münster, Germany; nedvetsk@uni-muenster.de

⁶ Residency Anaesthesiology, Department of Anaesthesiology und Critical Care Medicine, Technische Universität, Dresden, Germany


⁷ Facultad de Química, Universidad Autónoma de Querétaro, Mexico

⁸ Institut für Herz- und Kreislaufphysiologie Heinrich-Heine-Universität, Düsseldorf, Germany


*Corresponding Author: Prof Harald Schmidt, Department for Pharmacology and Personalised Medicine, MeHNS, FHML, Maastricht University, PO Box 616, 6200MD Maastricht, the Netherlands, Phone: +31-3-9905-5752; Email: hschmidt@ppmlab.net and melbatreek@ppmlab.net

Key words: endothelial cells; soluble guanylate cyclase; nitric oxide; cyclic guanosine mono phosphate; redox regulation; Bay 58-2667, apo-sGC


SUPPLEMENTARY FIGURES




Supplementary Fig. S1. 100 μM DETA-NO is the maximal concentration to downregulate sGC. Treatment of PPAECs for 72h with different concentrations of DETA-NO showed that 100 μM is the maximal concentration to downregulate both sGCa₁ (solid bars) and sGCβ₁ (open bars) (N=6). *P< 0.05 by one-way analysis of variance (ANOVA), data represent means ± SEM.



Supplementary Fig. S2. PPAECs display intact NO-cGMP-PKG signaling since treatment with 8-Br-cGMP (500 μM), or YC-1 for 72hrs reduced PKG expression (N=6-9). *P< 0.05 by one-way analysis of variance (ANOVA), data represent means ± SEM.



Supplementary Fig. S3. Inhibition of PKG does not prevent DETA-NO induced downregulation of sGCa₁ and sGC β₁. Treatment of PPAECs for 72h with either 100 μM DETA-NO, or with the PKG inhibitor Rp-8-Br-PET-cGMPS (10 μM), or a combination thereof resulted in down regulation of sGCa₁ (solid bars) and sGCβ₁ (open bars) level that could not be reversed by Rp-8-Br-PET-cGMPS (N=6). *P< 0.05 by one-way analysis of variance (ANOVA), data represent means ± SEM.



Supplementary Fig. S4. Representative Immune detection blots of controls, sGC α_1 and sGC β_1 . A. PPAECs. B. Mouse. C-E. ARDS (Multiple exposures).