

<b>Table 6: General recommendations for uveitis/rheumatologically patients on conventional IMT or biologics (n =139)</b>	
Hand and personal hygiene	139 (100%)
Avoiding crowds	139 (100%)
Use of masks	120 (86.3%)
Avoiding unproductive attendances at hospital	139 (100%)
Postponing long-term follow-up	127 (91.4%)
Self-isolation	107 (77%)
If really sick (with no symptoms of respiratory infection), see urgently doctor	136 (97.8%)

**Consensus for “Yes”**

**Table 6A:** General recommendations for uveitis patients on conventional IMT or biologics – As per **years of experience** and **number of patients of uveitis on IMT per year**

Criteria	Uveitis specialists' expertise		No. (%)
Hand and personal hygiene	Years of experience in uveitis	≤ 15 (n = 80)	80 ( 100 % )
		> 15 (n = 59)	59 ( 100 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	85 ( 100 % )
		≥ 300 (n =54)	54 ( 100 % )
Avoiding crowds	Years of experience in uveitis	≤ 15 (n = 80)	80 ( 100 % )
		> 15 (n = 59)	59 ( 100 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	85 ( 100 % )
		≥ 300 (n =54)	54 ( 100 % )
Use of masks	Years of experience in uveitis	≤ 15 (n = 80)	72 ( 90 % )
		> 15 (n = 59)	48 ( 81.36 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	74 ( 87.06 % )
		≥ 300 (n =54)	46 ( 85.19 % )
Avoiding unproductive attendances at hospital	Years of experience in uveitis	≤ 15 (n = 80)	80 ( 100 % )
		> 15 (n = 59)	59 ( 100 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	85 ( 100 % )
		≥ 300 (n =54)	54 ( 100 % )
Postponing long-term follow-up	Years of experience in uveitis	≤ 15 (n = 80)	76 ( 95 % )
		> 15 (n = 59)	51 ( 86.44 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	76 ( 89.41 % )
		≥ 300 (n =54)	51 ( 94.44 % )
Self-isolation	Years of experience in uveitis	≤ 15 (n = 80)	61 ( 76.25 % )
		> 15 (n = 59)	46 ( 77.97 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	64 ( 75.29 % )
		≥ 300 (n =54)	43 ( 79.63 % )
If really sick (with no symptoms of respiratory infection), see urgently doctor	Years of experience in uveitis	≤ 15 (n = 80)	78 ( 97.5 % )
		> 15 (n = 59)	58 ( 98.31 % )
	Uveitis patients on IMT/year	< 300 (n = 85)	82 ( 96.47 % )
		≥ 300 (n =54)	54 ( 100 % )

**Table 6B:** General recommendations for uveitis patients on conventional IMT or biologics -  
As per **geographical region of practice**

Criteria	Experts' geographical area	No. (%)
Hand and personal hygiene	Asia (n = 44)	44 ( 100 % )
	Africa (n = 3)	3 ( 100 % )
	Europe (n = 43)	43 ( 100 % )
	North America (n = 34)	34 ( 100 % )
	South America (n =7)	7 ( 100 % )
	Oceania (n = 8)	8 ( 100 % )
Avoiding crowds	Asia (n = 44)	44 ( 100 % )
	Africa (n = 3)	3 ( 100 % )
	Europe (n = 43)	43 ( 100 % )
	North America (n = 34)	34 ( 100 % )
	South America (n =7)	7 ( 100 % )
	Oceania (n = 8)	8 ( 100 % )
Use of masks	Asia (n = 44)	43 ( 97.73 % )
	Africa (n = 3)	3 ( 100 % )
	Europe (n = 43)	35 ( 81.4 % )
	North America (n = 34)	28 ( 82.35 % )
	South America (n =7)	7 ( 100 % )
	Oceania (n = 8)	4 ( 50 % )
Avoiding unproductive attendances at hospital	Asia (n = 44)	44 ( 100 % )
	Africa (n = 3)	3 ( 100 % )
	Europe (n = 43)	43 ( 100 % )
	North America (n = 34)	34 ( 100 % )
	South America (n =7)	7 ( 100 % )
	Oceania (n = 8)	8 ( 100 % )
Postponing long-term follow-up	Asia (n = 44)	38 ( 86.36 % )
	Africa (n = 3)	3 ( 100 % )
	Europe (n = 43)	42 ( 97.67 % )
	North America (n = 34)	30 ( 88.24 % )
	South America (n =7)	7 ( 100 % )
	Oceania (n = 8)	7 ( 87.5 % )
Self-isolation	Asia (n = 44)	34 ( 77.27 % )
	Africa (n = 3)	1 ( 33.33 % )
	Europe (n = 43)	35 ( 81.4 % )
	North America (n = 34)	24 ( 70.59 % )
	South America (n =7)	7 ( 100 % )
	Oceania (n = 8)	6 ( 75 % )
If really sick (with no symptoms of respiratory infection), see urgently doctor	Asia (n = 44)	43 ( 97.73 % )
	Africa (n = 3)	3 ( 100 % )
	Europe (n = 43)	43 ( 100 % )
	North America (n = 34)	33 ( 97.06 % )
	South America (n =7)	6 ( 85.71 % )
	Oceania (n = 8)	8 ( 100 % )

**Consensus for “Yes”**

