

Supporting Information

Repurposing Antiviral Drugs for Orthogonal RNA-Catalyzed Labeling of RNA

*Mohammad Ghaem Maghami, Surjendu Dey, Ann-Kathrin Lenz, and Claudia Höbartner**

anie_202001300_sm_miscellaneous_information.pdf

Table of Contents

1. Materials and DNA/RNA oligonucleotides	3
1.1. General materials	3
1.2. DNA oligonucleotides: Primers and transcription templates	3
1.3. RNA oligonucleotides: transcripts and synthetic RNAs	4
2. Experimental Procedures	6
2.1. Synthesis of tenofovir diphosphate analogs	6
2.2. In vitro selection of ribozymes using Biotin-tenofovir-DP	17
2.3. Characterization and application of FJ ribozymes	17
2.3.1 Single-turnover kinetic experiments	17
2.3.2 Preparative tenofovir labeling and probing of the modification site	18
2.3.3 Application of FJ ribozymes for labeling of cellular RNA	18
2.3.4 Mutually orthogonal dual-color labeling of a synthetic transcript	18
2.3.5. Probing of the modification site on the target cellular RNA using reverse transcription	19
3. Results and Discussion	20
3.1 Sequencing results of FJ ribozyme selection.....	20
3.1.1. FJ ribozyme sequences identified by Sanger sequencing.....	20
3.1.2. FJ ribozyme sequences identified by Illumina sequencing.....	20
3.2 Supporting Figures S1-S17	22
NMR spectra	33
References	59

SUPPORTING INFORMATION

1. Materials and DNA/RNA oligonucleotides

1.1. General materials

Unmodified oligonucleotides such as primers and transcription templates were ordered from Microsynth and subjected to PAGE purification as in our previous report.^[1]

Dynabeads streptavidin T1 and speedbead neutravidin coated-magnetic particles were purchased from Thermo Fisher scientific and GE-healthcare respectively. (Diethoxyphosphoryl)methyl 4-methylbenzenesulfonate, 6-FAM azide and Sulfo-Cy5 azide were purchased from Fluorochem and Jena Bioscience, respectively. All other chemicals were purchased from Sigma-Aldrich or ABCR and used without further purification. HPLC grade solvents were purchased from VWR.

Silica gel plates coated with fluorescent indicator were used for thin layer chromatography (TLC) and the plates were visualized with UV light. Silica gel 60, 0.032-0.063 mm (230-450 mesh) was used for column chromatography.

NMR spectra (¹H, ¹³C and ³¹P) were recorded using 400 MHz Bruker Avance III and Avance III HD spectrometers. Chemical shifts (¹H and ¹³C) were reported in parts per million (ppm) relative to TMS ($\delta = 0.00$ ppm) and were referenced to residual protium in the solvent. Coupling constants (*J*) are reported in Hz with the following multiplicity designations: s (singlet), d (doublet), t (triplet), q (quartet), doublet of doublet (dd), m (multiplet), and br (broad).

High-resolution ESI mass spectra in positive or negative ion mode were acquired on a Bruker micrOTOF-Q III.

Anion exchange chromatography was performed on a GE Healthcare ÄKTAprime plus system, on DEAE Sephadex A-25 (GE-Healthcare), self-packed 3x20 cm column. *Detection wavelength*: 280 nm, *Solvent systems*: buffer A: 100 mM TEAB (pH = 7.5); buffer B: 1.2 M TEAB (pH = 7.5), *Flow rate*: 6 ml/min, *Gradient*: 0 –100% buffer B in 200 min.

Fluorescent imaging of the kinetic and activity assay gels were taken using a BioRad Chemidoc gel-documentation device.

1.2. DNA oligonucleotides: Primers and transcription templates

Description	5' -Sequence-3'
T7 Promoter	CTGTAATACGACTCACTATA
Forward primer/2 nd PCR fwd primer	TTGAAGGCTCAGTATGTCTTATAGTGAGTCGTATTACA
Pool reverse primer	GGTAAGGTGGACATACTG-N40-GCCTTCAAGGATGGTAGGCTGG
Selection reverse primer	GGTAAGGTGGACATACTG
1 st PCR forward primer	CTTCAACCAGCCTACCATCC
Cloning forward primer	TAAATAAAATAACTGTAATACGACTCACTATAGGACATACTGAGC
Fwd primer for substrate sequence deletion	CTGTAATACGACTCACTATAGGCCAGCCTACCATCC
E. coli 5S rRNA forward primer	CTGTAATACGACTCACTATAGGTGCCTGGCGCCGTAGCGCGGTGGTCCCACCTGACCCCATGCCG AACTCAGAAGTGAA
E. coli 5S rRNA reverse primer	TGCCTGGCAGTTCCCTACTCTCGCATGGGGAGACCCACACTACCATCGGGCTACGGCGTTTCAC TTCTGAGTTCGG
1st PCR NGS N8 UMI forward primer	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGNNNNNNNAACCAGCCTACCATCC
1st PCR NGS reverse primer	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGAAGGTCGACATACTG
Indexed forward primer I5_S502	AATGATACGGCGACCACCGAGATCTACACCTCTCTATTTCGTCGGCAGCGTC
Indexed reverse primer I7_N703 (Round 12)	CAAGCAGAAGACGGCATAACGAGATTTCTGCCTGTCTCGTGGGCTCGG
Indexed reverse primer I7_N704 (Round 7)	CAAGCAGAAGACGGCATAACGAGATGCTCAGGAGTCTCGTGGGCTCGG
RT primer 16S 344-361 (A325)	CCCCTGCTGCCTCCCGT
RT primer 16S 413-430 (A383)	TACAACCCGAAGGCCTTC
RT primer 16S 687-704 (A649)	TCTACGCATTTCCACCGCT
RT primer 23S 322-339 (A272)	ACGGGGCTGTACACCTGT
RT primer 23S 653-672 (A637)	GGTCTATACCCTGCAACT
RT primer 23S 1599-1618 (A1572)	TGTGTCGGTTTGGGGTACGA
ssDNA templates for	5' -Sequence-3'
Parent substrate sequence	TTGAAGGCTCAGTATGTCTTATAGTGAGTCGTATTACAG
TM	CCAGGAACCTCGACGCACCCCTATAGTGAGTCGTATTACAG
TV1	AACTTCCCTCTCATAACCTATAGTGAGTCGTATTACAG
TV2	GGTCCTTCTCCTGCGTGCCCTATAGTGAGTCGTATTACAG
Parent-AAG	TTGAAGGCTTAGTATGTCTTATAGTGAGTCGTATTACAG
Parent -UAG	TTGAAGGCTAAGTATGTCTTATAGTGAGTCGTATTACAG
Parent -CAG	TTGAAGGCTGAGTATGTCTTATAGTGAGTCGTATTACAG
Parent -GAA	TTGAAGGTTTTCAGTATGTCTTATAGTGAGTCGTATTACAG

SUPPORTING INFORMATION

Parent -GAU	TTGAAGGATCAGTATGTCCTATAGTGAGTCGTATTACAG
Parent -GAC	TTGAAGGGTCAAGTATGTCCTATAGTGAGTCGTATTACAG
TM-UGAGC	CCAGGAGCTCAACGCACCCTATAGTGAGTCGTATTACAG
5'TV1-3'parent	TTGAAGGCTCTCATACACCTATAGTGAGTCGTATTACAG
TV1-GAGC	AACTTCGCTCTCATACACCTATAGTGAGTCGTATTACAG
RNA w two labeling sites (DLRNA)	CATTAACCTCGACGCATCCTTAATTGAAGGCTCAGTATGTCCTATAGTGAGTCGTATTACAG
ssDNA templates for	5' -Sequence-3'
FJ1	GGACATACTGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTTCAACCTATAGTGAGTCGTATTACAG
FJ8	GGACATACTGGGGGAAGGTAATGTCCTAAGGTTTCGTTTTGTAAACACCGTGCCTTCAACCTATAGTGAGTCGTATTACAG
FJC1	GGACATACTGGGAACTTGCTAGTTTACCCTTTTCTTATGTGACACGTTTCGCCTTCAACCTATAGTGAGTCGTATTACAG
FJC3	GGACATACTGGAAATTTGGAGGCGGCAATATACGGTTCGGTGGCATTCGGCCTTCAACCTATAGTGAGTCGTATTACAG
FJC9	GGACATACTGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGCCTTCAACCTATAGTGAGTCGTATTACAG
FJ1_TM	GGGTGCGTCGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGTTCTTGGCCTATAGTGAGTCGTATTACAG
FJ1_TV1	GGTGTATGAGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGGGAAGTTCCTATAGTGAGTCGTATTACAG
FJ1_TV2	GGCAGCAGGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGAAGACCCCTATAGTGAGTCGTATTACAG
FJ1_TM-UGAGC	GGGTGCGTTGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTCCTGGCCTATAGTGAGTCGTATTACAG
FJ1_5'TV1-3'Par	GGTGTATGAGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTTCAACCTATAGTGAGTCGTATTACAG
FJ1_TV1-GAGC	GGTGTATGAGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCGAAGTTCCTATAGTGAGTCGTATTACAG
FJ1_AAG	GGACATACTATAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTTCAACCTATAGTGAGTCGTATTACAG
FJ1_GAA	GGACATACTGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGACCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_TM	GGGTGCGTCGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGTTCCTGGCCTATAGTGAGTCGTATTACAG
FJC9_TV1	GGTGTATGAGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGGGAAGTTCCTATAGTGAGTCGTATTACAG
FJC9_TV2	GGCAGCAGGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGAAGGACCCCTATAGTGAGTCGTATTACAG
FJC9_AAG	GGACATACTAGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGCCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_UAG	GGACATACTTGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGCCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_CAG	GGACATACTCGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGCCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_GAA	GGACATACTGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTACCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_GAU	GGACATACTGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTTCCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_GAC	GGACATACTGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTCCCTTCAACCTATAGTGAGTCGTATTACAG
FJC9_5S-A99	TCCCCATGCGGGATGCCCAATTCGCTAAAGTGTAGTGCAACATATCTCGTGAGTAGGGAACTTATAGTGAGTCGTATTAC
FJ1_23S-A272	AGGAGCCAGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTGAATCCTATAGTGAGTCGTATTACAG
FJ1_23S-A362	ACATGCTGTGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTCGATGCCTATAGTGAGTCGTATTACAG
FJ1_23S-A1572	TTCCAGGAAATAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTTCAACCTATAGTGAGTCGTATTACAG
FJ1_16S-A383	AATGGGCGCATAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTGATGCAGCCTATAGTGAGTCGTATTAC
FJ1_16S-A649	GATACTGGCATAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTTGAGTCCTATAGTGAGTCGTATTAC
FJ1_16S-A860	GTGGCTTCCGGTAGACCCTTGCCAAAGGATCTTCAGTTTTATGAGGGTGGGCTAACGCCTCCTATAGTGAGTCGTATTAC
FH14_16S-A325	ACTGGAAGTGTCTATGTTTGCAGCGTTTGCAGTATGTCTACTCAGCGTGGACACGGTCCCTATAGTGAGTCGTATTACAG
FH14_16S-A412	GCGTGTATGATCTATGTTTGCAGCGTTTGCAGTATGTCTACTCAGCGTGGAGGCCTCCTATAGTGAGTCGTATTACAG
FH14_23S-A272	AGGAGCCAGTCTATGTTTGCAGCGTTTGCAGTATGTCTACTCAGCGTGGCCTGAATCCTATAGTGAGTCGTATTACAG
FH14_23S-A637	AGGAAACCGTCTATGTTTGCAGCGTTTGCAGTATGTCTACTCAGCGTGGTCTAACCTATAGTGAGTCGTATTACAG
FH14_23S-A1572	TTCCAGGAAATCTATGTTTGCAGCGTTTGCAGTATGTCTACTCAGCGTGGCCTTCAACCTATAGTGAGTCGTATTACAG
FH14_A34_DLRNA	GGATGCGTCGTCTATGTTTGCAGCGTTTGCAGTATGTCTACTCAGCGTGGTTAATGCCTATAGTGAGTCGTATTACAG

1.3. RNA oligonucleotides: transcripts and synthetic RNAs

Pool and substrate sequences (modification site shown in bold)

Description	5' -Sequence-3'
Selection pool	GGACAUACUGAGCCUUAACCAGCCUACCAUCCUUGAAGGC-N40-CAGUAUGUCCACCUUACC
Parent substrate sequence	GGACAUACUG A GCCUUCAA
TM	GGGUGCGUCC A GUUCCUGG
TV1	GGUGUAUGAG A GGGAAGUU
TV2	GGCAGCAGG A GAAGGACC
TM-UGAGC	GGGUGCGUUG A GCUCCUGG
5'TV1-3'parent	GGUGUAUGAG A GCCUUCAA
TV1-GAGC	GGUGUAUGAG A GCGAAGUU
Parent-AAG	GGACAUACUAAGCCUUCAA
Parent -UAG	GGACAUACUU A GCCUUCAA
Parent -CAG	GGACAUACUC A GCCUUCAA
Parent -GAA	GGACAUACUG A ACCUUCAA
Parent -GAU	GGACAUACUG A UCCUUCAA
Parent -GAC	GGACAUACUG A CCCUUCAA
3'-aminohexyl RNA substrate	ACAUACUG A GCCUUCAA-C6-NH2
5'-Hexynyl RNA substrate	Hexyne-GACAUACUG A GCCUUCAAAUA

SUPPORTING INFORMATION

RNA w two labeling sites (DLRNA)	GGACAUACUG <u>AG</u> CCUCAAUUAAGGAUGCGUCG <u>AG</u> UUAUG
E. coli 5S rRNA (In vitro transcribed)	GGUGCCUGGCGGCCGUAAGCGCGGUGGCCACCUGACCCCAUGCCGAACUCAGAAGUGA AACGCCGUAAGCGCCGAUGGUAGUGUGGGGUCUCCCCAUGCGAG <u>AG</u> UAGGGGAACUGCCAG GCAU



Ribozymes (Recognition arms underlined)

Name	5'-Sequence-3'
FJ1	GGUUGAAGGCCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCUA <u>CAGUAUGUCC</u>
FJ8	GGUUGAAGGCCGAAACGUGUCACAUAAAACGGUAAAACUAGCAAGUUC <u>CAGUAUGUCC</u>
FJC1	GGUUGAAGGCCGAAUGCCACCGAACCGUAUAAUUGCCGCCUCCAAUUC <u>CAGUAUGUCC</u>
FJC3	GGUUGAAGGCCACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CAGUAUGUCC</u>
FJC9	GGCCAGGA <u>ACC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCUA <u>CGACGCACCC</u>
FJ1_TM	GGAA <u>CUUCC</u> CCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCUA <u>CUCAUACACC</u>
FJ1_TV1	GGGGUCCU <u>UCC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCUA <u>CCUGCGUGCC</u>
FJ1_TV2	GGAA <u>CUUCC</u> ACCGGUGUUAACAAAACGAAACCUUAGGACAUUACCUUCC <u>CCUCAUACACC</u>
FJ1_TM-UGAGC	GGUUGAAGGCCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>ACUCAUACACC</u>
FJ1_5'TV1-3'par	GGAA <u>CUUCC</u> ACCCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>ACUCAUACACC</u>
FJ1_TV1-GAGC	GGUUGAAGGCCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCUA <u>UAGUAUGUCC</u>
FJ1_AAG	GGUUGAAGGU <u>CC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>ACAGUAUGUCC</u>
FJ1_GAA	GGUUGAAGGCACGGUGUUAACAAAACGAAACCUUAGGACAUUACCUUCC <u>CCUCAUACACC</u>
FJC9_TM	GGAA <u>CUUCC</u> ACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CCUCAUACACC</u>
FJC9_TV1	GGGGUCCU <u>UCC</u> ACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CCUGCGUGCC</u>
FJC9_TV2	GGUUGAAGGCCACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>UAGUAUGUCC</u>
FJC9_AAG	GGUUGAAGGCCACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>AAAGUAUGUCC</u>
FJC9_UAG	GGUUGAAGGCCACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>GAGUAUGUCC</u>
FJC9_CAG	GGUUGAAGGU <u>AC</u> GAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CCAGUAUGUCC</u>
FJC9_GAA	GGUUGAAGGAACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CAGUAUGUCC</u>
FJC9_GAU	GGUUGAAGGGACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CAGUAUGUCC</u>
FJC9_GAC	GGAUUCAGGCCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>ACUGGGCUCCU</u>
FJC9_5S-A99	GGU <u>UCCUACUC</u> ACGAGAUUUGUACACUUUAGCGAAUUGGGCAUCC <u>CCGCAUGGGGA</u>
FJ1_23S-A272	GGCA <u>UCGAGCC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>ACACAGCAUGU</u>
FJ1_23S-A362	GGU <u>UAGAGGCC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>AUUUCCUGGAA</u>
FJ1_23S-A1572	GG <u>AUCAAGGCC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>CAAUGUUCAG</u>
FJ1_16S-A383	GG <u>ACUC</u> AAGGCCACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>UAGCCAGUAUC</u>
FJ1_16S-A649	GGACCG <u>GUUAGCC</u> ACCCUCAUAAAACUGAAGAUCUUUGGCAAGGGUCU <u>ACCGGAAGCCAC</u>
FJ1_16S-A860	GG <u>ACCGU</u> UCCACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGACAGU <u>UCCAGU</u>
FH14_16S-A325	GGAGGCCU <u>UCC</u> ACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGAU <u>CAUACACGC</u>
FH14_16S-A412	GGUGCGGG <u>ACC</u> ACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGAU <u>AACCCAACA</u>
FH14_23S-A272	GG <u>AUUC</u> AGGCCACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGACUGGG <u>CUCCU</u>
FH14_23S-A637	GGUUAAGACCACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGACGGU <u>UCCCU</u>
FH14_23S-A1572	GGUUAAGAGGCCACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGAU <u>UCCUGGAA</u>
FH14_A34-DLRNA	GGCAUUAACCACGCUGAGUAGACAUACUUGCAAACGCUGCAAACAUAGACGACGCA <u>UCC</u>

SUPPORTING INFORMATION

2. Experimental Procedures


2.1. Synthesis of tenofovir diphosphate analogs

Scheme S1. Synthesis of *N*⁶-biotin-tenofovir diphosphate (biotin-Ten-DP).Scheme S2. a) Synthesis of tenofovir diphosphate, *N*⁶-modified analogs *N*⁶-hexynyl-Ten-DP and *N*⁶-azidoheptyl-Ten-DP. b) Synthesis of fluorescently labeled tenofovir diphosphates by CuAAC.

SUPPORTING INFORMATION

List of abbreviations used in the synthetic procedures:

ACN = Acetonitrile, CDI = 1,1'-Carbonyldiimidazole, Boc = *tert*-Butyloxycarbonyl, Bu₃N = Tri-*n*-butylamine, Cy5 = sulfo-cyanine-5, DCM = Dichloromethane, DMF = Dimethylformamide, DP = diphosphate, FAM = Fluorescein, h = hours, min = minutes, NHS = *N*-Hydroxysuccinimide, o.n. = overnight, RP = reversed-phase, RT = room temperature, TEAA = Triethylammonium acetate, TEAB = Triethylammonium bicarbonate, Ten = tenofovir, TLC = Thin-layer chromatography, TMSBr = Bromotrimethylsilane, Ts = Tosyl.

1-(6-chloro-9H-purin-9-yl)propan-2-one (1)


Compound **1** was synthesized following a published procedure with slight modifications.^[2]

To a solution of 6-chloropurine (1 g, 6.47 mmol) in 30 ml dry DMF, anhydrous K₂CO₃ (894 mg, 6.47 mmol) was added. The mixture was stirred for 30 min at RT, followed by addition of a solution of chloroacetone (0.57 ml, 7.12 mmol) in 10 ml dry DMF over 1.5 h. The reaction mixture was continuously stirred overnight at RT, then filtered through Celite, and the filtrate was evaporated to dryness. The residue was dissolved in 150 ml CHCl₃ and washed with saturated NaHCO₃ (4 x 25 ml) and brine (20 ml). The organic layer was dried over Na₂SO₄, and the solvent was evaporated. The crude product was purified by flash column chromatography (3% MeOH/DCM), affording compound **1** as a yellowish-white solid (804 mg, 3.82 mmol, 59%). A minor amount of 7-alkylated isomer was formed, which was separated by column chromatography.

¹H NMR (400 MHz, CDCl₃): δ 8.71 (s, 1H), 8.11 (s, 1H), 5.15 (s, 2H), 2.37 (s, 3H).

¹³C NMR (101 MHz, CDCl₃): δ 198.6, 152.1, 151.8, 151.2, 145.6, 131.1, 52.3, 27.2.

HRMS (ESI⁺): Exact mass calculated for C₈H₈ClN₄O [M + H]⁺, 211.03811. Found 211.03798.


1-(6-chloro-9H-purin-9-yl)propan-2-ol (2)

Compound **1** (780 mg, 3.7 mmol) was dissolved in 40 ml MeOH/ACN (2:1). The solution was cooled to 0 °C and NaBH₄ (168 mg, 4.44 mmol) was added in three portions over 20 min. The mixture was stirred for another 20 min at 0 °C until TLC confirmed the complete consumption of the reactant. Excess of NaBH₄ reagent was consumed by the addition of 20 ml saturated NH₄Cl and the reaction volume was concentrated to ~25 ml. The reaction mixture was extracted with CHCl₃ (4 x 50 ml). The combined organic layers were washed with saturated NH₄Cl (30 ml), brine (30 ml), dried over Na₂SO₄, and concentrated under reduced pressure. The crude product was purified by flash column chromatography (4-5% MeOH/DCM), to give the pure title compound **2** as a white solid (731 mg, 3.44 mmol, 93%).

¹H NMR (400 MHz, CDCl₃): δ 8.70 (s, 1H), 8.21 (s, 1H), 4.43 (dd, *J* = 14.0, 2.7 Hz, 1H), 4.31 (dq, *J* = 8.0, 6.3, 2.7 Hz, 1H), 4.13 (dd, *J* = 14.1, 8.0 Hz, 1H), 3.13 (s, 1H), 1.32 (d, *J* = 6.3 Hz, 3H).

¹³C NMR (101 MHz, CDCl₃): δ 151.8, 151.7, 150.6, 146.4, 131.0, 66.0, 51.5, 20.8.

HRMS (ESI⁺): Exact mass calculated for C₈H₉ClN₄NaO [M + Na]⁺, 235.03571. Found 235.03518.

Diethyl (((1-(6-chloro-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonate (3)

SUPPORTING INFORMATION

In a 25 ml Schlenk flask, a colorless solution of compound **2** (150 mg, 0.71 mmol) in 5 ml dry DMF was treated with $\text{Mg}(\text{O}^i\text{Bu})_2$ (363 mg, 2.13 mmol) at RT. The white suspension was stirred at 70 °C for 30 min. Then (diethoxyphosphoryl)methyl 4-methylbenzenesulfonate (344 mg, 1.07 mmol) was added in small portions over 10 min under argon flow. The reaction mixture was then stirred for 7 h at 70 °C. The solvent was evaporated to dryness, and the residue was put onto a pad of celite and washed with 20% MeOH/DCM. The filtrate was reduced in vacuo and the crude residue was purified by flash chromatography (1.5-3.5% MeOH/DCM) to afford compound **3** as a colorless oil (225 mg, 0.62 mmol, 87%).


¹H NMR (400 MHz, CDCl_3): δ 8.73 (s, 1H), 8.31 (s, 1H), 4.47 (dd, $J = 14.5, 2.8$ Hz, 1H), 4.22 (dd, $J = 14.5, 7.7$ Hz, 1H), 4.11 – 4.05 (m, 2H), 4.04 – 3.98 (m, 2H), 3.97 – 3.91 (m, 1H), 3.85 (dd, $J = 13.7, 8.9$ Hz, 1H), 3.58 (dd, $J = 13.7, 9.4$ Hz, 1H), 1.32 – 1.29 (t, $J = 7.1$ Hz, 3H), 1.26 – 1.20 (m, 6H).

¹³C NMR (101 MHz, CDCl_3): δ 152.0, 151.8, 150.9, 146.5, 131.2, 75.9 (d, $J = 11.2$ Hz), 63.5, 62.4 (d, $J = 6.5$ Hz), 62.3 (d, $J = 6.6$ Hz), 61.8, 48.6, 16.5 – 16.4 (m).

³¹P NMR (162 MHz, CDCl_3): δ 20.66.

HRMS (ESI⁺): Exact mass calculated for $\text{C}_{13}\text{H}_{21}\text{ClN}_4\text{O}_4\text{P}$ [$\text{M} + \text{H}$]⁺, 363.09835. Found 363.09836.

tert-butyl (6-((9-(2-((diethoxyphosphoryl)methoxy)propyl)-9H-purin-6-yl)amino)hexyl)carbamate (4)



In a 50 ml Schlenk flask, a solution of compound **3** (205 mg, 0.57 mmol) and *N*-Boc-1,6-hexanediamine hydrochloride (358 mg, 1.41 mmol) in 12 ml absolute EtOH was treated with *N,N*-diisopropylethylamine (0.48 ml, 2.83 mmol). The solution was heated under reflux for 18 h until TLC confirmed the complete consumption of the reactants. The solvent was evaporated and the residue was purified by flash column chromatography (4-4.5% MeOH/DCM). The pure title compound **4** was obtained as a colorless sticky oil (246 mg, 0.45 mmol, 80%).


¹H NMR (400 MHz, MeOD): δ 8.24 (s, 1H), 8.07 (s, 1H), 4.35 (dd, $J = 14.5, 3.1$ Hz, 1H), 4.20 (dd, $J = 14.5, 7.9$ Hz, 1H), 4.06 – 3.90 (m, 6H), 3.71 (dd, $J = 14.0, 9.7$ Hz, 1H), 3.58 (br s, 2H), 3.03 (t, $J = 6.8$ Hz, 2H), 1.70 (p, $J = 7.1$ Hz, 2H), 1.50 – 1.36 (m, 15H), 1.27 – 1.18 (m, 9H).

¹³C NMR (101 MHz, MeOD): δ 158.5, 156.1, 153.7, 149.8, 142.9, 120.0, 79.7, 77.6 (d, $J = 12.3$ Hz), 64.0 (d, $J = 6.6$ Hz), 63.9 (d, $J = 6.6$ Hz), 63.8, 62.1, 49.1, 41.5, 41.3, 30.9, 30.5, 28.8, 27.7, 27.6, 16.7 – 16.6 (m).

³¹P NMR (162 MHz, MeOD): δ 21.99.

HRMS (ESI⁺): Exact mass calculated for $\text{C}_{24}\text{H}_{44}\text{N}_6\text{O}_6\text{P}$ [$\text{M} + \text{H}$]⁺, 543.30545. Found 543.30534.

((1-(6-((6-aminohexyl)amino)-9H-purin-9-yl)propan-2-yl)oxy)methylphosphonic acid (5)



Deprotection of the phosphonate ethyl ester and Boc groups was performed according to modified published procedures.^{[3],[4]}

In a 25 ml Schlenk flask, compound **4** (162 mg, 0.3 mmol) was dissolved in 6 ml dry DCM. The tube was placed in an ice bath, and TMSBr (0.24 ml, 1.8 mmol) was added dropwise to the solution. The reaction mixture was stirred for 3 h at 0 °C until TLC confirmed the complete consumption of compound **4**. Evaporation of the solvent afforded a yellowish sticky oil, which was co-evaporated with MeOH followed by Et_2O . The obtained yellowish-white solid was dissolved in 20 ml of water and extracted with DCM (4 x 3 ml). The aqueous fraction was lyophilized, and the title compound **5** was obtained as a white solid in quantitative yield.

¹H NMR (400 MHz, D_2O): δ 8.34 (s, 1H), 8.30 (s, 1H), 4.44 (dd, $J = 14.9, 3.0$ Hz, 1H), 4.26 (dd, $J = 14.8, 7.8$ Hz, 1H), 4.00 – 3.94 (m, 1H), 3.77 (dd, $J = 13.8, 8.8$ Hz, 1H), 3.57 – 3.51 (m, 2H), 2.94 (t, $J = 7.6$ Hz, 2H), 1.77 – 1.68 (m, 2H), 1.66 – 1.58 (m, 2H), 1.49 – 1.34 (m, 4H), 1.19 (d, $J = 6.3$ Hz, 3H).

¹³C NMR (101 MHz, D_2O): δ 148.5, 147.0, 144.7, 144.1, 118.0, 76.5 (d, $J = 11.6$ Hz), 63.5 (d, $J = 161.0$ Hz), 48.4, 42.1, 39.3, 27.2, 26.5, 25.3, 25.2, 15.8.


SUPPORTING INFORMATION

³¹P NMR (162 MHz, D₂O): δ 19.04.

HRMS (ESI⁻): Exact mass calculated for C₁₅H₂₆N₆O₄P [M - H]⁻, 385.17586. Found 385.17591.

HRMS (ESI⁺): Exact mass calculated for C₁₅H₂₈N₆O₄P [M + H]⁺, 387.19042. Found 387.18994.

(((1-(6-((6-(5-((3a*S*,4*S*,6a*R*)-2-oxohexahydro-1*H*-thieno[3,4-*d*]imidazole-4-yl)pentanamido)exyl)amino)-9*H*-purin-9-yl)propan-2-yl)oxy)methyl)phosphonic acid (Biotin-tenofovir, **6)**



In a 10 ml Schlenk flask, compound **5** (40 mg, 0.104 mmol) was dissolved in 3 ml dry DMF. Biotin-NHS ester (42 mg, 0.124 mmol), was added and the reaction mixture was stirred for 15 min at RT before Et₃N (150 μl, 1.08 mmol) was added. The white suspension became clear after stirring for 30 min at RT, and the reaction progress was monitored by RP-TLC (using 20% MeCN/H₂O). The reaction mixture was stirred overnight at RT before the solvent was evaporated. After coevaporation with H₂O, the residue was dissolved in 30 ml water and extracted with DCM (3 x 3 ml). The aqueous fraction was lyophilized, and a yellowish-white solid was obtained. The crude product was purified by reversed-phase chromatography to give the title compound **6** as a white foam (40 mg, 0.065 mmol, 62%).


RP-purification procedure: *Performed on*: GE Healthcare ÄKTAprime plus, *Column*: Lobar 310-25 LiChroprep RP-18 (40-63 μm), Merck *Detection wavelength*: 280 nm, *Solvent systems*: A: H₂O, B: ACN, *Gradient*: 5–50% B in 50 min, *Flow rate*: 7 ml/min

¹H NMR (400 MHz, CD₃CN:D₂O (1:1)): δ 8.78 (s, 2H), 5.03 – 5.00 (m, 1H), 4.91 (dd, *J* = 14.7, 3.6 Hz, 1H), 4.84 (dd, *J* = 7.9, 4.5 Hz, 1H), 4.48 – 4.41 (m, 1H), 4.19 (dd, *J* = 12.9, 9.3 Hz, 1H), 4.10 – 3.99 (m, 2H), 3.75 – 3.64 (m, 6H), 3.42 (dd, *J* = 12.9, 5.0 Hz, 1H), 3.22 (d, *J* = 12.9 Hz, 1H), 2.70 (t, *J* = 7.3 Hz, 2H), 2.56 – 2.54 (m, 1H), 2.25 – 1.85 (m, 14H), 1.78 (t, *J* = 7.3 Hz, 4H), 1.63 (d, *J* = 6.3 Hz, 3H).

³¹P NMR (162 MHz, CD₃CN:D₂O (1:1)): δ 15.70.

HRMS (ESI⁻): Exact mass calculated for C₂₅H₄₀N₈O₆PS [M - H]⁻, 611.25346. Found 611.25471.

(((1-(6-((6-(5-((3a*S*,4*S*,6a*R*)-2-oxohexahydro-1*H*-thieno[3,4-*d*]imidazole-4-yl)pentanamido)hexyl)amino)-9*H*-purin-9-yl)propan-2-yl)oxy)methyl-[hydroxy(phosphonoxy)phosphoryl]oxy-phosphinic acid, sodium salt (Biotin-tenofovir-DP; **7)**



Phosphonate activation using CDI, followed by reaction with pyrophosphate was performed in analogy to published procedures.^[5]

First, the tributylammonium salt of compound **6** was generated. A suspension of the free acid form of **6** (21 mg, 34 μmol) in 1 ml of dry DMF was treated with tributylamine (32 μl, 136 μmol) for 1 h at RT. The clear solution was evaporated, followed by co-evaporation with dry DMF (3 x 1 ml). Then, the tributylammonium salt of compound **6** was dissolved in 1 ml dry DMF, and a solution of CDI (28 mg, 170 μmol) in 0.5 ml of dry DMF was added dropwise. The reaction mixture was stirred for 18 h at RT. Excess CDI was quenched by the addition of MeOH (5.5 μl, 136 μmol) followed by stirring for 1 h at RT. Then, (Bu₃NH)₂H₂P₂O₇ (93 mg, 170 μmol) dissolved in 1 ml dry DMF was added. The resulting white suspension was stirred overnight at RT. The supernatant was separated by centrifugation and the precipitate (imidazolium pyrophosphate) was washed with dry DMF (2 x 0.2 ml). The combined DMF fractions were evaporated, followed by co-evaporation with H₂O. The resulting residue was dissolved in 10 ml H₂O and extracted with DCM (3 x 2 ml). The aqueous fraction was lyophilized, and the crude product was purified by anion exchange chromatography (on DEAE Sephadex, see general

SUPPORTING INFORMATION

methods, section 1.1). The product-containing fraction (55-65% buffer B) was evaporated, and then co-evaporated with H₂O. The residue was further purified by reversed-phase chromatography. The resulting triethylammonium salt of the title compound was obtained as a white foam, and was transformed to the sodium salt by precipitation from 20 ml 2% NaClO₄ in acetone. After centrifugation, the pellet was washed with acetone (3 x 5 ml), and dried in vacuo. The pure product **7** was obtained as a white solid (5.5 mg, 6.39 μmol, 19%).

RP-purification procedure: *Column*: Lobar 310-25 LiChroprep RP-18 (40-63 μm), Merck

Detection wavelength: 280 nm, *Solvent systems*: buffer A: 100 mM TEAB (pH = 7.5); buffer B: 100 mM TEAB in 80% ACN


Gradient: 10% B for 10 min, 10-25% B in 15 min, 25% B for 5 min, 25-35% B in 10 min, 35% B for 10 min, *Flow rate*: 5 ml/min

¹H NMR (400 MHz, D₂O): δ 8.25 (s, 2H), 4.47 – 4.42 (m, 2H), 4.30 – 4.25 (m, 2H), 4.05 – 3.99 (m, 1H), 3.87 – 3.68 (m, 2H), 3.54 (q, *J* = 7.1 Hz, 4H), 3.20 – 3.12 (m, 3H), 2.83 (dd, *J* = 13.1, 5.0 Hz, 1H), 2.63 (d, *J* = 13.0 Hz, 1H), 2.22 – 2.17 (m, 2H), 1.71 – 1.26 (m, 14H), 1.17 – 1.12 (m, 7H).

³¹P NMR (162 MHz, D₂O): δ 8.72 (d, *J* = 26.4 Hz), -10.70 (d, *J* = 19.9 Hz), -23.11 (t, *J* = 21.5 Hz).

HRMS (ESI): Exact mass calculated for C₂₅H₄₂N₈O₁₂P₃S [M - H]⁻, 771.18612. Found 771.18521.

Diethyl (((1-(6-amino-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonate (Tenofovir diethyl ester, **8**)



In a stainless steel autoclave, compound **3** (200 mg, 0.55 mmol) was treated with 25 ml of 7 N NH₃ in MeOH for 2 days at 50 °C. After solvent evaporation, the residue was directly purified by flash column chromatography (3-7.5% MeOH/DCM) to give the desired compound **8** as a white solid (**8**, 112 mg, 0.326 mmol, 59%).

¹H NMR (400 MHz, CDCl₃): δ 8.33 (s, 1H), 7.96 (s, 1H), 6.00 (br s, 2H), 4.35 (dd, *J* = 14.4, 3.0 Hz, 1H), 4.15 – 3.99 (m, 5H), 3.95 – 3.88 (m, 1H), 3.83 (dd, *J* = 13.6, 9.1 Hz, 1H), 3.58 (dd, *J* = 13.6, 9.7 Hz, 1H), 1.29 (t, *J* = 7.1 Hz, 3H), 1.26 – 1.22 (m, 6H).

¹³C NMR (101 MHz, CDCl₃): δ 155.5, 152.8, 150.1, 141.8, 119.1, 76.3 (d, *J* = 11.9 Hz), 63.6, 62.4 (d, *J* = 6.7 Hz), 62.3 (d, *J* = 6.7 Hz), 61.9, 48.2, 16.4 (d, *J* = 5.2 Hz), 16.3 (d, *J* = 5.2 Hz).

³¹P NMR (162 MHz, CDCl₃): δ 20.83.

HRMS (ESI⁺): Exact mass calculated for C₁₃H₂₃N₅O₄P [M + H]⁺, 344.14822. Found 344.14823.

The MeO-substituted compound **8b** was isolated as a side product (eluted first from the column, colorless oil, 68 mg, 0.190 mmol, 35%)


¹H NMR (400 MHz, CDCl₃): δ 8.51 (s, 1H), 8.07 (s, 1H), 4.40 (dd, *J* = 14.5, 3.0 Hz, 1H), 4.20 – 3.99 (m, 8H), 3.96 – 3.89 (m, 1H), 3.82 (dd, *J* = 13.7, 9.1 Hz, 1H), 3.57 (dd, *J* = 13.6, 9.6 Hz, 1H), 1.30 (t, *J* = 7.1 Hz, 3H), 1.28 – 1.20 (m, 6H).

¹³C NMR (101 MHz, CDCl₃): δ 161.0, 152.1, 151.9, 143.4, 121.1, 76.3 (d, *J* = 11.9 Hz), 63.5, 62.4 (d, *J* = 6.5 Hz), 62.3 (d, *J* = 6.5 Hz), 61.9, 54.2, 48.3, 16.5 – 16.3 (m).

³¹P NMR (162 MHz, CDCl₃): δ 20.72.

HRMS (ESI⁺): Exact mass calculated for C₁₄H₂₄N₄O₅P [M + H]⁺, 359.14788. Found 359.14783.

(((1-(6-amino-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonic acid (Tenofovir, **9**)



Unmodified Tenofovir was synthesized in analogy to previous reports.^[3, 6]

In a 25 ml Schlenk flask, compound **8** (100 mg, 0.29 mmol) was dissolved in 6 ml dry DCM and treated with TMSBr (0.23 ml, 1.74 mmol) at 0 °C for 3 h. TLC confirmed the complete consumption of the reactant. The solvent was evaporated, and the resulting yellowish sticky oil was co-evaporated with MeOH, followed by Et₂O. The residue was dissolved in 20 ml of water and washed with DCM (4 x 3 ml). The aqueous fraction was lyophilized, and the crude product was purified by silica gel column chromatography (acetone/H₂O/Et₃N,

SUPPORTING INFORMATION

6:1:1), followed by a second purification by silica gel column chromatography (*i*PrOH/H₂O/conc. NH₃, 7:2:1) to afford compound **9** as a white foam (63 mg, 0.22 mmol, 75%).


¹H NMR (400 MHz, D₂O): δ 8.14 (s, 1H), 8.02 (s, 1H), 4.25 (dd, *J* = 14.6, 3.9 Hz, 1H), 4.14 (dd, *J* = 14.6, 5.8 Hz, 1H), 3.92 – 3.84 (m, 1H), 3.50 (dd, *J* = 12.6, 9.3 Hz, 1H), 3.37 (dd, *J* = 12.6, 9.3 Hz, 1H), 1.03 (d, *J* = 6.3 Hz, 3H).

¹³C NMR (101 MHz, D₂O): δ 155.1, 151.9, 148.8, 143.2, 117.7, 75.4 (d, *J* = 11.3 Hz), 66.2 (d, *J* = 153.0 Hz), 47.4, 16.1.

³¹P NMR (162 MHz, D₂O): δ 13.92.

HRMS (ESI): Exact mass calculated for C₉H₁₃N₅O₄P [M - H]⁻, 286.07106. Found 286.07109.

[1-(6-amino-9H-purin-9-yl)propan-2-yl]oxymethyl-[hydroxy(phosphonoxy)phosphoryl]oxy-phosphinic acid, triethylammonium salt (Tenofovir-DP, **10)**



Compound **9** (25 mg, 86 μmol) was co-evaporated with dry DMF (3 x 1 ml), dissolved in 1 ml of dry DMF and treated with tributylamine (82 μl, 344 μmol) for 1 h at RT. To the clear solution, a solution of CDI (105 mg, 648 μmol) in 1 ml dry DMF was added dropwise, and the reaction mixture was stirred for 18 h at RT. Excess CDI was quenched by the addition of MeOH (22 μl, 557 μmol). After 1 h at RT, (Bu₃NH)₂H₂P₂O₇ (354 mg, 645 μmol) dissolved in 1.5 ml dry DMF was added, and the resulting white suspension was stirred overnight at RT. The precipitate was removed by centrifugation and the supernatant was evaporated. The residue was directly loaded onto an anion exchange column (see general method section). Two fractions were collected separately (first fraction: 60-70% buffer B, second fraction: 95-100% buffer B), and after evaporation further purified via reversed-phase chromatography. Compound **10** was obtained from the first eluting fraction as white foam (21.4 mg, 25 μmol, 29%). The side product from the second fraction was identified as the homo-coupled side product (5.8 mg, 3.5 μmol, 4%).

RP-purification procedure: *Performed on:* GE Healthcare ÄKTAprime plus, *Column:* Lobar 310-25 LiChroprep RP-18 (40-63 μm), Merck *Detection wavelength:* 280 nm, *Solvent systems:* buffer A: 100 mM TEAB (pH = 7.5); buffer B: 100 mM TEAB in 80% ACN *Gradient:* 5% B for 5 min, 5-15% B in 15 min, 15% B for 25 min, *Flow rate:* 5 ml/min

Compound **10**:

λ_{max} (UV-vis absorption in H₂O): 259 nm

¹H NMR (400 MHz, D₂O): δ 8.32 (s, 1H), 8.23 (s, 1H), 4.42 (dd, *J* = 14.7, 3.3 Hz, 1H), 4.24 (dd, *J* = 14.8, 5.9 Hz, 1H), 4.04 – 3.97 (m, 1H), 3.84 (dd, *J* = 13.3, 9.2 Hz, 1H), 3.74 (dd, *J* = 13.3, 9.5 Hz, 1H), 1.09 (d, *J* = 6.3 Hz, 3H).

³¹P NMR (162 MHz, D₂O): δ 8.67 (d, *J* = 26.4 Hz), -10.93 (d, *J* = 19.2 Hz), -23.35 (t, *J* = 22.3 Hz).

HRMS (ESI): Exact mass calculated for C₉H₁₅N₅O₁₀P₃ [M - H]⁻, 446.00373. Found 446.00372.

A second fraction was obtained from anion exchange chromatography, as a white foam after RP-purification which was identified as the homo-coupled side product (5.8 mg, 3.5 μmol, 4%).


λ_{max} (UV-vis absorption in H₂O): 292 nm

¹H NMR (400 MHz, D₂O): δ 8.92 (s, 1H), 8.82 (s, 1H), 4.61 (dd, *J* = 14.7, 2.9 Hz, 1H), 4.42 (dd, *J* = 14.8, 6.2 Hz, 1H), 4.12 – 4.05 (m, 1H), 3.89 (dd, *J* = 13.4, 9.1 Hz, 1H), 3.75 (dd, *J* = 13.2, 9.7 Hz, 1H), 1.19 (d, *J* = 6.3 Hz, 3H).

³¹P NMR (162 MHz, D₂O): δ 8.70 (d, *J* = 26.5 Hz), -11.00 (d, *J* = 19.9 Hz), -23.45 (t, *J* = 22.6 Hz).

HRMS (ESI): Exact mass calculated for C₁₉H₂₈N₁₀O₂₁P₆ [M - 2H]²⁻, 458.99336. Found 458.99559.

Diethyl (((1-(6-(hex-5-yn-1-ylamino)-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonate (Hexynyl-tenofovir diethyl ester, **11)**



SUPPORTING INFORMATION

In a 50 ml Schlenk flask, compound **3** (200 mg, 0.55 mmol) and 5-Hexyn-1-amine (134 mg, 1.38 mmol) were dissolved in 15 ml absolute EtOH and treated with *N,N*-diisopropylethylamine (0.24 ml, 1.40 mmol). The solution was heated under reflux for 18 h. TLC confirmed the complete consumption of the reactant. The solvent was evaporated and the residue was purified by flash chromatography (3.5% MeOH/DCM). The pure product **11** was obtained as a colorless sticky oil (221 mg, 0.52 mmol, 95%).


¹H NMR (400 MHz, CDCl₃): δ 8.36 (s, 1H), 7.88 (s, 1H), 5.76 (br s, 1H), 4.33 (dd, *J* = 14.4, 2.9 Hz, 1H), 4.14 – 4.00 (m, 5H), 3.95 – 3.88 (m, 1H), 3.81 (dd, *J* = 13.6, 9.1 Hz, 1H), 3.68 (br s, 2H), 3.57 (dd, *J* = 13.8, 9.4 Hz, 1H), 2.25 (td, *J* = 7.0, 2.6 Hz, 2H), 1.95 (t, *J* = 2.9 Hz, 1H), 1.85 – 1.77 (m, 2H), 1.69 – 1.62 (m, 2H), 1.30 (t, *J* = 7.1 Hz, 3H), 1.27 – 1.22 (m, 6H).

¹³C NMR (101 MHz, CDCl₃): δ 154.9, 153.0, 149.0, 140.9, 119.3, 83.9, 76.4 (d, *J* = 12.0 Hz), 68.7, 63.6, 62.4 (d, *J* = 6.5 Hz), 62.3 (d, *J* = 6.5 Hz), 61.9, 48.1, 40.0, 28.8, 25.6, 18.1, 16.5 (d, *J* = 6.8 Hz), 16.4 (d, *J* = 5.4 Hz).

³¹P NMR (162 MHz, CDCl₃): δ 20.84.

HRMS (ESI⁺): Exact mass calculated for C₁₉H₃₁N₅O₄P [M + H]⁺, 424.21082. Found 424.21083.

(((1-(6-(hex-5-yn-1-ylamino)-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonic acid (Hexynyl-tenofovir, 12)



In a 25 ml Schlenk flask, compound **11** (100 mg, 0.24 mmol) was dissolved in 6 ml dry DCM, and TMSBr (0.20 ml, 1.51 mmol) was added dropwise at 0 °C. The reaction was stirred for 3 h at 0 °C. TLC confirmed the complete consumption of the reactant. The solvent was evaporated and the resulting residue was co-evaporated with MeOH and Et₂O before it was dissolved in 20 ml water and extracted with DCM (4 x 3 ml). The aqueous fraction was lyophilized and purified by silica gel column chromatography (iPrOH/H₂O/conc. NH₃, 7:2:1) to afford compound **12** as a white foam (78 mg, 0.22 mmol, 90%).


¹H NMR (400 MHz, D₂O): δ 8.08 (s, 1H), 8.05 (s, 1H), 4.26 (dd, *J* = 14.6, 3.4 Hz, 1H), 4.12 (dd, *J* = 14.7, 6.6 Hz, 1H), 3.92 – 3.84 (m, 1H), 3.61 (dd, *J* = 13.1, 9.3 Hz, 1H), 3.46 – 3.40 (m, 3H), 2.26 (t, *J* = 2.6 Hz, 1H), 2.16 (td, *J* = 7.0, 2.6 Hz, 2H), 1.72 – 1.65 (m, 2H), 1.56 – 1.49 (m, 2H), 1.09 (d, *J* = 6.3 Hz, 3H).

¹³C NMR (101 MHz, D₂O): δ 153.1, 150.7, 147.4, 142.7, 117.9, 85.5, 75.8 (d, *J* = 11.9 Hz), 69.3, 64.8 (d, *J* = 157.8 Hz), 47.7, 40.4, 27.5, 25.0, 17.3, 16.0.

³¹P NMR (162 MHz, D₂O): δ 15.73.

HRMS (ESI⁻): Exact mass calculated for C₁₅H₂₁N₅O₄P [M - H]⁻, 366.13366. Found 366.13366.

[1-(6-(hex-5-yn-1-ylamino)-9H-purin-9-yl)propan-2-yl]oxymethyl-[hydroxy(phosphonoxy)phosphoryl]oxy-phosphinic acid, triethylammonium salt (Hexynyl-tenofovir-DP, 13)



The free acid form of compound **12** (65 mg, 177 μmol) was suspended in 1 ml dry DMF and treated with tributylamine (168 μl, 708 μmol) for 1 h at RT to form a clear solution of the corresponding tributylammonium salt. The solvent was evaporated, followed by co-evaporation with dry DMF (3 x 1 ml). The tributylammonium salt of compound **12** was dissolved in 1.5 ml dry DMF and a solution of CDI (216 mg, 1.33 mmol) in 1.5 ml dry DMF was added dropwise. The reaction mixture was stirred for 18 h at RT before excess CDI was quenched by the addition of MeOH (46 μl, 1.15 mmol) and stirring was continued for 1 h at RT. Then (Bu₃NH)₂H₂P₂O₇ (730 mg, 1.33 mmol) dissolved in 2 ml dry DMF was added and the resulting white suspension was stirred overnight at RT. The precipitate was removed by centrifugation and the supernatant was evaporated, followed by co-evaporation with H₂O. The residue was dissolved in 10 ml H₂O and extracted with DCM (3 x 2 ml). The aqueous layer was lyophilized and the crude product was purified by anion exchange chromatography (see general method section). The product-containing fraction (60-70% buffer B) was collected and the solvent was evaporated. The final purification was achieved by reverse-phase chromatography to obtain compound **13** as a white foam (35.5 mg, 38 μmol, 21%).

SUPPORTING INFORMATION


RP-purification procedure: *Performed on:* GE Healthcare ÄKTApurime plus, *Column:* Lobar 310-25 LiChroprep RP-18 (40-63 μm), Merck
Detection wavelength: 280 nm, *Solvent systems:* buffer A: 100 mM TEAB (pH = 7.5); buffer B: 100 mM TEAB in 80% ACN
Gradient: 5% B for 5 min, 5-20% B in 5 min, 20% B for 10 min, 20-30% B in 5 min, 30% B for 20 min, *Flow rate:* 5 ml/min

^1H NMR (400 MHz, D_2O): δ 8.21 (s, 1H), 8.20 (s, 1H), 4.40 (dd, J = 14.8, 3.5 Hz, 1H), 4.25 (dd, J = 14.8, 5.7 Hz, 1H), 4.06 – 3.99 (m, 1H), 3.84 (dd, J = 13.3, 9.2 Hz, 1H), 3.74 (dd, J = 13.3, 9.4 Hz, 1H), 3.58 (br s, 2H), 2.29 (t, J = 2.6 Hz, 1H), 2.24 (td, J = 7.0, 2.6 Hz, 2H), 1.82 – 1.75 (m, 2H), 1.65 – 1.58 (m, 2H), 1.11 (d, J = 6.3 Hz, 3H).

^{31}P NMR (162 MHz, D_2O): δ 8.64 (d, J = 26.6 Hz), -10.92 (d, J = 19.9 Hz), -23.36 (dd, J = 26.5, 19.9 Hz).

HRMS (ESI $^-$): Exact mass calculated for $\text{C}_{15}\text{H}_{23}\text{N}_5\text{O}_{10}\text{P}_3$ [$\text{M} - \text{H}$] $^-$, 526.06633. Found 526.06663.

Diethyl (((1-(6-((6-azidohexyl)amino)-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonate (Azidoheptyl-tenofovir diethyl ester, **14)**



In a 50 ml Schlenk flask, compound **3** (150 mg, 0.41 mmol) and 6-azidoheptan-1-amine **21** (150 mg, 1.05 mmol) were dissolved in 5 ml absolute EtOH and treated with *N,N*-diisopropylethylamine (0.18 ml, 1.05 mmol). Then the solution was refluxed for 18 h. TLC confirmed the complete consumption of the reactant. The solvent was evaporated and the residue was purified by flash chromatography (3.5% MeOH/DCM) to afford the pure product **14** as yellowish sticky oil (185 mg, 0.39 mmol, 95%).

^1H NMR (400 MHz, CDCl_3): δ 8.36 (s, 1H), 7.88 (s, 1H), 5.77 (s, 1H), 4.32 (dd, J = 14.4, 3.0 Hz, 1H), 4.13 – 4.00 (m, 5H), 3.95 – 3.88 (m, 1H), 3.81 (dd, J = 13.6, 9.1 Hz, 1H), 3.64 – 3.54 (m, 3H), 3.25 (t, J = 6.9 Hz, 2H), 1.73 – 1.66 (m, 2H), 1.63 – 1.56 (m, 2H), 1.48 – 1.40 (m, 4H), 1.29 (t, J = 7.1 Hz, 3H), 1.27 – 1.22 (m, 6H).

^{13}C NMR (101 MHz, CDCl_3): δ 154.7, 152.9, 148.9, 140.7, 119.0, 76.2 (d, J = 12.0 Hz), 63.5, 62.2 (d, J = 6.7 Hz), 62.1 (d, J = 6.7 Hz), 61.8, 51.1, 47.9, 40.4, 29.4, 28.6, 26.3, 16.3 (d, J = 8.1 Hz), 16.2 (d, J = 5.3 Hz).

^{31}P NMR (162 MHz, CDCl_3): δ 20.84.

HRMS (ESI $^+$): Exact mass calculated for $\text{C}_{19}\text{H}_{34}\text{N}_8\text{O}_4\text{P}$ [$\text{M} + \text{H}$] $^+$, 469.24351. Found 469.24351.

(((1-(6-((6-azidoheptyl)amino)-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonic acid (Azidoheptyl-tenofovir, **15)**



In a 25 ml Schlenk flask, compound **14** (100 mg, 0.21 mmol) was dissolved in 4 ml dry DCM and the solution was cooled in an ice bath. TMSBr (0.20 ml, 1.51 mmol) was added dropwise and the reaction mixture was stirred for 3 h at 0 °C. TLC confirmed the complete consumption of the reactant. The solvent was evaporated, co-evaporated with MeOH and Et_2O , before the resulting yellowish-white solid was dissolved in 20 ml water. After washing with DCM (4 x 3 ml), the aqueous fraction was lyophilized to afford compound **15** as a white solid in quantitative yield.

^1H NMR (400 MHz, D_2O): δ 8.24 (s, 1H), 8.22 (s, 1H), 4.32 (dd, J = 14.6, 3.0 Hz, 1H), 4.13 (dd, J = 14.8, 7.8 Hz, 1H), 3.87 – 3.79 (m, 1H), 3.68 (dd, J = 13.8, 8.8 Hz, 1H), 3.48 – 3.41 (m, 3H), 3.10 (t, J = 6.8 Hz, 2H), 1.65 – 1.50 (m, 2H), 1.42 – 1.32 (m, 2H), 1.31 – 1.13 (m, 4H), 1.06 (d, J = 6.3 Hz, 3H).


^{13}C NMR (101 MHz, D_2O): δ 148.2, 146.8, 144.6, 144.3, 117.4, 76.41 (d, J = 11.8 Hz), 63.08 (d, J = 161.8 Hz), 50.9, 48.5, 43.9, 42.2, 27.7, 27.3, 25.5, 15.8.

^{31}P NMR (162 MHz, D_2O): δ 19.65.

HRMS (ESI $^-$): Exact mass calculated for $\text{C}_{15}\text{H}_{24}\text{N}_8\text{O}_4\text{P}$ [$\text{M} - \text{H}$] $^-$, 411.16636. Found 411.16638.

SUPPORTING INFORMATION

[1-(6-((6-azidohexyl)amino)-9H-purin-9-yl)propan-2-yl]oxymethyl-[hydroxy(phosphonooxy)phosphoryl]oxy-phosphinic acid, triethylammonium salt (Azidoheyl-tenofovir-DP, 16)



The free acid form of compound **15** (55 mg, 134 μ mol) was suspended in 1 ml dry DMF and treated with tributylamine (128 μ l, 536 μ mol) for 1 h at RT. The clear solution of the resulting tributylammonium salt was evaporated, and dried by further co-evaporation with dry DMF (3 x 1 ml), before it was dissolved in 1.5 ml dry DMF. A solution of CDI (109 mg, 670 μ mol) in 1.5 ml dry DMF was added dropwise, and the reaction mixture was stirred for 6 h at RT. Excess CDI was quenched by the addition of MeOH (22 μ l, 536 μ mol). After 1 h at RT, a solution of (Bu₃NH)₂H₂P₂O₇ (550 mg, 1 mmol) in 2 ml dry DMF was added, and the resulting white suspension was stirred overnight at RT. The supernatant was collected and separated from the precipitate (imidazolium pyrophosphate) by centrifugation. The solvent was evaporated, the residue was dissolved in 10 ml H₂O and extracted with DCM (3 x 2 ml). The aqueous solution was lyophilized and the crude product was purified by anion exchange chromatography (see general method section). The product-containing fraction (60-70% buffer B) was collected, the solvent was evaporated and the residue was subjected to purification by RP chromatography. The product **16** was obtained as a white foam (42.3 mg, 43 μ mol, 32%).


RP-purification procedure: *Performed on:* GE Healthcare ÄKTAprime plus, *Column:* Lobar 310-25 LiChroprep RP-18 (40-63 μ m), Merck *Detection wavelength:* 280 nm, *Solvent systems:* buffer A: 100 mM TEAB (pH = 7.5); buffer B: 100 mM TEAB in 80% ACN *Gradient:* 10-40% B in 30 min, 40% B for 10 min, 40-50% B in 10 min, 50% B for 10 min, *Flow rate:* 5 ml/min

¹H NMR (400 MHz, D₂O): δ 8.21 (s, 1H), 8.19 (s, 1H), 4.40 (dd, J = 14.7, 3.6 Hz, 1H), 4.25 (dd, J = 14.7, 5.6 Hz, 1H), 4.06 – 3.99 (m, 1H), 3.84 (dd, J = 13.3, 9.3 Hz, 1H), 3.74 (dd, J = 13.4, 9.3 Hz, 1H), 3.56 (br s, 2H), 3.27 (t, J = 6.8 Hz, 2H), 1.73 – 1.64 (m, 2H), 1.60 – 1.54 (m, 2H), 1.46 – 1.35 (m, 4H), 1.11 (d, J = 6.3 Hz, 3H).

³¹P NMR (162 MHz, D₂O): δ 8.66 (d, J = 26.2 Hz), -10.36 (d, J = 20.2 Hz), -23.10 (t, J = 22.6 Hz).

HRMS (ESI): Exact mass calculated for C₁₅H₂₆N₈O₁₀P₃ [M - H]⁻, 571.09902. Found 571.09905.

FAM-tenofovir-DP (17)



In a small 1.5 mL reaction tube 500 nmol compound **13** (10 μ l of 50 mM in DMF; 1 eq) were combined with 10 μ l 6-FAM azide (100 mM in DMF, 2 eq) and 10 μ l DMF. In another 1.5 mL reaction tube, 5 μ l CuSO₄ (100 mM in H₂O; 1 eq) was mixed with 15 μ l of a freshly prepared solution of sodium ascorbate (200 mM in H₂O; 6 eq), and added to the first 1.5 mL reaction tube, to give a final volume of 50 μ l (final DMF/H₂O 3:2). The reaction mixture was stirred overnight at RT. Then, 1.2 ml ice-cold 2% NaClO₄ in acetone was added and centrifuged (centrifuge at 4 $^{\circ}$ C, 15000 rpm for 30 min). The precipitate was dissolved in water and purified by reversed phase HPLC. The isolated product was lyophilized to obtain compound **17** as a red foam (100 nmol, 20%), which was dissolved in 100 μ l water to prepare a 1 mM stock solution for further use.


RP-purification procedure: *Performed on:* GE Healthcare ÄKTAmicro, *Column:* Nucleosil 100-5 C18 column, 250 x 4.6 mm, Macherey-Nagel, *Detection wavelength:* 260 nm and 496 nm, *Solvent systems:* buffer A: 100 mM TEAA (pH = 7.0); buffer B: 100 mM TEAA in 80% CAN, *Gradient:* 5% B for 2.5 min, 5-20% B in 5 min, 20-22.5% B in 10 min, 22.5-50% B in 2.5 min, 50% B for 2.5 min, 50-100% B in 2.5 min, *Column oven temperature:* 40 $^{\circ}$ C, *Flow rate:* 1 ml/min

HRMS (ESI): Exact mass calculated for C₃₉H₄₁N₉O₁₆P₃ [M - H]⁻, 984.18896. Found 984.18897.

Exact mass calculated for C₃₉H₄₀N₉O₁₆P₃ [M - 2H]²⁻, 491.59084. Found 491.59459.

SUPPORTING INFORMATION

Cy5-tenofovir-DP (18)




In a small 1.5 mL reaction tube 500 nmol compound **13** (10 μ L of 50 mM in H₂O; 1 eq) were combined with 10 μ L Sulfo-Cy5 azide (100 mM in H₂O, 2 eq) and 60 μ L DMF. In another 1.5 mL reaction tube, 5 μ L CuSO₄ (100 mM in H₂O; 1 eq) were mixed with 15 μ L of a freshly prepared solution of sodium ascorbate (200 mM in H₂O; 6 eq), and the mixture was added to the first 1.5 mL reaction tube, to give a final volume of 100 μ L (final DMF/H₂O 3:2). A small magnetic stir bar was put into the 1.5 mL reaction tube and the mixture was stirred overnight at RT. The solution was split into two 1.5 mL reaction tubes (2 x 50 μ L) and 1.2 ml ice-cold 2% NaClO₄ in acetone was added to each tube. After centrifugation (at 4 °C, 15000 rpm for 30 min), the precipitate was dissolved in water and purified by reversed phase HPLC. The compound **18** was obtained as a dark blue foam (120 nmol, 24%), which was dissolved in 120 μ L water to make a 1 mM stock concentration for further use.

RP-purification procedure: *Performed on:* GE Healthcare ÄKTAmicro, *Column:* Nucleosil 100-5 C18 column, 250 x 4.6 mm, Macherey-Nagel, *Detection wavelength:* 260 nm and 647 nm, *Solvent systems:* buffer A: 100 mM TEAA (pH = 7.0); buffer B: 100 mM TEAA in 80% CAN, *Gradient:* 5% B for 2.5 min, 5-20% B in 2.5 min, 20-40% B in 22.5 min, 40-100% B in 2.5 min, *Column oven temperature:* 40 °C, *Flow rate:* 1 ml/min

HRMS (ESI): Exact mass calculated for C₅₂H₇₀N₁₁O₂₀P₃S₃ [M - 2H]²⁻, 678.65923. Found 678.65925.

Exact mass calculated for C₅₂H₆₉N₁₁O₂₀P₃S₃ [M - 3H]³⁻, 452.10373. Found 452.10752.

2-(6-bromohexyl)isoindoline-1,3-dione (19)



Compound **19** was synthesized according to a modified published procedure.^[7]


In a 250 ml Schlenk flask, 1,6-Dibromohexane (15.6 ml, 101.41 mmol) was dissolved in 75 ml dry DMF, and potassium phthalimide (8 g, 43.19 mmol) was added in small portions over 30 min. The reaction mixture was stirred overnight at RT, and then the solvent was evaporated. The white residue was put onto celite and washed with 200 ml EtOAc. The filtrate was evaporated and the crude product was purified by flash chromatography (15% EtOAc/Hexane), affording compound **19** as a white solid (9.72 g, 31.34 mmol, 73%).

¹H NMR (400 MHz, CDCl₃): δ 7.84 – 7.82 (m, 2H), 7.72 – 7.69 (m, 2H), 3.70 – 3.66 (m, 2H), 3.39 (t, *J* = 6.8 Hz, 2H), 1.88 – 1.81 (m, 2H), 1.72 – 1.65 (m, 2H), 1.51 – 1.44 (m, 2H), 1.40 – 1.32 (m, 2H).

¹³C NMR (101 MHz, CDCl₃): δ 168.4, 133.9, 132.1, 123.2, 37.8, 33.7, 32.6, 28.3, 27.7, 26.0.

HRMS (ESI*): Exact mass calculated for C₁₄H₁₇BrNO₂ [M + H]⁺, 310.04372. Found 310.04457.

SUPPORTING INFORMATION

2-(6-azidohexyl)isoindoline-1,3-dione (**20**)


Compound **20** was synthesized in analogy to a published procedure.^[7]

In a 100 ml round-bottom flask, compound **19** (9 g, 29.01 mmol) was dissolved in 30 ml DMF, followed by the addition of NaN_3 (5.66 g, 87.03 mmol) in portions over 1 h at RT. The reaction mixture was stirred overnight at RT. Then the solvent was evaporated and 150 ml DCM was added to the residue, followed by the extraction with water (4 x 50 ml) and brine (50 ml). The organic layer was dried over Na_2SO_4 , and after evaporation of the solvent, the title compound **20** was obtained as colorless sticky liquid which turned solid (7.58 g, 27.84 mmol, 96%) on cooling with ice water. Compound **20** was used for the next reaction without further purification.

^1H NMR (400 MHz, CDCl_3): δ 7.85 – 7.81 (m, 2H), 7.72 – 7.68 (m, 2H), 3.69 – 3.66 (m, 2H), 3.24 (t, J = 6.9 Hz, 2H), 1.72 – 1.64 (m, 2H), 1.62 – 1.55 (m, 2H), 1.45 – 1.32 (m, 4H).

^{13}C NMR (101 MHz, CDCl_3): δ 168.4, 133.9, 132.1, 123.1, 51.3, 37.8, 28.6, 28.4, 26.3, 26.2.

HRMS (ESI⁺): Exact mass calculated for $\text{C}_{14}\text{H}_{16}\text{N}_4\text{NaO}_2$ [$\text{M} + \text{Na}$]⁺, 295.11655. Found 295.11656.

6-azidohexan-1-amine (**21**)

Compound **21** was synthesized according to a modified published procedure.^[7]

In a 100 ml round-bottom flask, compound **20** (4 g, 14.69 mmol) was dissolved in 20 ml EtOH. To that solution, hydrazine hydrate (3.7 g) was added dropwise and the reaction was stirred overnight at RT. The reaction mixture turned to a white solid, which was crashed into small parts, followed by the addition of 75 ml EtOAc. After stirring for 1 h at RT, the reaction mixture was filtered and the white residue was washed with 200 ml EtOAc. The filtrate was evaporated and the remaining (yellowish liquid) was redissolved in 50 ml EtOAc, and treated with 50% aq. HCl. The clear solution turned cloudy after acidification. Then the acidic EtOAc solution was extracted with H_2O (4 x 15 ml). Aqueous fractions were combined and basified with concentrated aq. NaOH solution until the formation of liquid droplets was observed. The basified aq. solution was extracted with DCM (4 x 100 ml). The organic layers were combined and again extracted with H_2O (50 ml) followed by brine (50 ml). The DCM solution was dried over Na_2SO_4 , followed by solvent evaporation to yield compound **21** as yellowish liquid (1.54 g, 10.83 mmol, 74%).

^1H NMR (400 MHz, CDCl_3): δ 3.24 (t, J = 6.9 Hz, 2H), 2.67 (t, J = 7.0 Hz, 2H), 1.61 – 1.54 (m, 2H), 1.46 – 1.30 (m, 7H).

^{13}C NMR (101 MHz, CDCl_3): δ 51.4, 42.1, 33.6, 28.8, 26.6, 26.5.

HRMS (ESI⁺): Exact mass calculated for $\text{C}_6\text{H}_{15}\text{N}_4$ [$\text{M} + \text{H}$]⁺, 143.12912. Found 143.12917.

SUPPORTING INFORMATION

2.2. In vitro selection of ribozymes using Biotin-tenofovir-DP

The in vitro selection was performed in analogy to our previously described procedure for identification of FH ribozymes.^[1] The conditions for selection of FJ ribozymes are summarized as follows: In the first round of selection, 3 nmol of the N40 pool mixed with ~150 pmol of the 3'-lucifer yellow labeled N40 pool was dissolved in 60 μ L of the selection buffer (50 mM HEPES, pH 7.5, 120 mM KCl, 5 mM NaCl) including 40 mM MgCl₂ and 300 μ M of Biotin-Tenofovir-diphosphate (**7**). Following incubation overnight at 37°C, the unreacted biotin-tenofovir substrate was removed using ethanol precipitation. The reactive species were then immobilized on 1 mg of neutravidin beads (pre-treated with *E. coli* tRNA), and the bulk of the unreactive species in the pool were washed away using denaturing wash buffer. The beads were then suspended in formamide elution buffer and placed at 95°C for 7 min. The supernatant of this step was then subjected to ethanol precipitation. The eluted RNA was reverse transcribed and subjected to 10 rounds of PCR amplification. The primers in this PCR step do not encompass the modification site, therefore providing equal amplification opportunity to the modified RNA as the unmodified inactive carry-over RNA. A 2nd PCR was then performed (18-30 rounds) with an alternative primer which restores the substrate sequence to its full length and adds the T7 promoter. The product of the 2nd PCR was then transcribed in vitro with T7 RNA polymerase to generate the RNA pool entering the next selection round.

In all the subsequent selection rounds the pool concentration was maintained at ~60 μ M, including ~150 pmol of 3'-LY labeled RNA pool of the same round. In the following selection rounds the amount of RNA pool and the reaction volume used was adjusted based on the yield of the previous round and amount of beads used in the capture step was kept at 0.1 mg of the beads per 100 pmol of RNA pool. To prevent selection of neutravidin binders from the pool, the affinity matrix was alternated between neutravidin and streptavidin every two rounds. The frequency of the active variants was estimated by comparing the fluorescence intensity of the eluted RNA at the end of the round to that of the total pool prior to the capture step. First sign of enrichment became apparent at the end of the 4th selection round where ~0.44% of the pool was retained. A rise in the activity level to ~1.42% in the next round further confirmed the success of the enrichment process. The incubation time was reduced at round 8 from overnight to 4 h which was further reduced to 1 h by the 12th round. A streptavidin gel shift assay on the outcome of the round 12 showed over 80% activity after overnight incubation. The pool also demonstrated trans-activity when the substrate sequence was removed from the pool and added as a separate entity to the reaction. At this point the pool was cloned (using a TOPO-TA cloning kit) and 10 representative colonies were subjected to Sanger sequencing. Two unique sequences were identified which were denoted as FJ1 and FJ8.

The pool was then further examined for additional variants by subjecting it to Next Generation Sequencing (NGS). To prepare the NGS libraries, RT-PCR samples from round 7 and round 12 were subjected to two consecutive PCR amplifications during which the unimolecular identifiers (UMIs) and indices were added. The sample from the last PCR was agarose gel purified and submitted to the Core Unit Systems Medicine (Uni Würzburg) for amplicon sequencing (NextSeq-500 HighOutput 75nt single end). New sequence variants were identified in the demultiplexed and trimmed fastq files (see below, section 3.1.2), three of which (denoted as FJC1, FJC3 and FJC9) were chosen for further characterization.

2.3. Characterization and application of FJ ribozymes

2.3.1 Single-turnover kinetic experiments

Kinetic experiments were performed as previously reported.^[1] Briefly, a 10:1 ratio of the ribozyme : fluorescently labeled substrate sequence were dissolved in selection buffer resulting in 10 μ M and 1 μ M final concentrations of ribozyme and the substrate sequence (i.e. single-turnover conditions). The reactions were performed in the presence of 40 mM MgCl₂ and 300 μ M of the Tenofovir-diphosphate analog substrate. The reaction was incubated at 37°C and 1 μ L aliquots were taken from the experiment at desired time-points and quenched by adding to 4 μ L of stop solution. Half of the resulting samples were resolved on 20% analytical denaturing PAGE for 1 h under constant power of 25 W for 25 cm plates and 35 W for 30 cm plates. For kinetic experiments in which different tenofovir analogs were tested the running time was increased to 1.5 h to achieve proper resolution. The gels were imaged on a Chemidoc device and band intensities were quantified by ImageLab software. The fraction of labeled product was calculated, and the timecourse fit to single-exponential kinetics according to $Y = Y_{\max}(1 - \exp(-k_{\text{obs}}t))$.

2.3.1.1 Target sequence requirement of FJ ribozymes

To investigate target sequence generality of FJ ribozymes, transition or transversion mutations were introduced on the original parent substrate sequence, initially outside the GAG modification context. FJ ribozymes were also designed with complementary arms to these sequences. Single turn-over kinetic experiments were then performed to determine the capacity of the ribozymes for modifying broader range of target sequences. Since FJ1 and FJ8 ribozymes were unable to modify the extensively mutated substrate sequence variants, a new set of mutants were designed to further investigate their modification context requirement. In one of the mutants the unchanged nucleotides was expanded by one nucleotide from each side of the GAG context, while the rest of the sequence was subjected to transition mutations, resulting in TM-UGAGC, in which UGAGC is identical to the parent substrate sequence. Another sequence was also designed in which the segment 5' to the GAG was transversion mutated while the 3'-segment was kept unchanged resulting in 5'TV1-3'parent sequence. Single-turnover kinetic experiments using these substrates determined the C nucleotide following the GAG context is essential. Further investigations determined the minimal modification context as 5'-AGC-3'. Optimal modification efficiency was obtained with sequences containing 5'-RAGCY-3' as the modification site context sequence.

SUPPORTING INFORMATION

2.3.1.2 Tenofovir-Diphosphate analog scope of FJ ribozymes

To determine nucleotide analog specificity of FJ ribozymes, unmodified tenofovir-diphosphate, *N*⁶-hexynyl-tenofovir-diphosphate and *N*⁶-azidoethyl-tenofovir diphosphate were employed in single turnover kinetic assays. None of the tested variants showed any dependency to the biotin moiety or the C6 linker as all the tested derivatives including the unmodified tenofovir-DP were efficiently ligated to the substrate sequence.

2.3.2. Preparative tenofovir labeling and probing of the modification site**2.3.2.1 Preparative modification of the substrate sequence with biotin-tenofovir**

One nmol of 5'-fluorescein labeled synthetic substrate sequence was mixed with 1.3 nmol of the FJ1 ribozyme in a total volume of 25 μ L of selection buffer including 40 mM MgCl₂ and 500 μ M of the *N*⁶-biotin-TenDP substrate. The sample was incubated at 37°C overnight, followed by quenching via addition of 25 μ L of high-dye solution. The sample was then resolved on a 20% denaturing PAGE and the band corresponding to the modified RNA was excised from the gel. The modified RNA product was then extracted from the gel and ethanol precipitated resulting in an isolated yield of 55%.

2.3.2.2 Probing of the modification site

Alkaline hydrolysis and T1 probing was used to locate the exact position of the modification site in the substrate sequence. For alkaline hydrolysis, 10 pmol of the modified or unmodified 5'-fluorescently labeled synthetic RNA substrate was incubated in 10 μ L of 50, 25 or 10 mM NaOH and incubated at 95°C. Two μ L aliquots were taken from each reaction after 1, 2.5 and 5 min of incubation and quenched by adding to 3 μ L of stop solution and placing on ice. For T1 digestion, 10 pmol of the modified or unmodified RNA substrate were dissolved in 5 μ L of 50 mM Tris pH= 7.5 containing 0.1 U/ μ L of RNase T1. The sample was placed at 37°C and quenched after 30 sec by adding 5 μ L of stop solution and placing on ice.

2.5 μ L of each of the timepoint samples and T1 probing samples were resolved on a 20% analytical denaturing PAGE for 1 h under constant power of 35 W and subjected to fluorescent imaging.

2.3.2.3. Debranching assay

5 pmol of 5'-fluorescein labeled substrate sequence modified using FJ1 or FH14 was dissolved in 5 μ L of Dbr1 reaction buffer (50 mM Tris pH = 7.4, 25 mM NaCl, 2.5 mM DTT, 0.01% (V/V) Tween-20, 0.15% (V/V) glycerol) including 1 or 5 μ M MnCl₂ and 0.5 ng/ μ L of recombinant Dbr1 (kindly provided by A. Hoskins, Univ. of Wisconsin). The sample was incubated at 37°C and 1 μ L timepoint samples were taken at 0, 15, 30 and 60 min and quenched by adding to 4 μ L of stop solution and placing in liquid nitrogen. 2.5 μ L of each of these samples were then resolved on a 20% analytical denaturing PAGE for 1 h under constant power of 25W. The gel was then subjected to fluorescent imaging.

2.3.3 Application of FJ ribozymes for labeling of cellular RNA

Total cellular RNA from TOP10 E.coli cells was isolated as previously described.^[1, 8]

For the experiments in which 16S and/or 23S rRNA were labeled, 20 ng of total cellular RNA from Top10 E. Coli cells (NEB) was mixed with 50 pmol of the corresponding ribozyme(s). The reaction was performed in 5 μ L of the selection buffer including 40 mM MgCl₂ and 200 μ M of the fluorescently labeled TenDP and/or ATP analogs. After 6h incubation at 37°C, the reaction was subjected to 2x isopropanol precipitation. The resulting pellet was washed using ice cold 70% ethanol. The dried pellet was dissolved in 2 μ L of stop solution and 1 μ L of milliQ water. The sample was then resolved on 1.3% agarose gel under 90 V for 45 min. The gel was then subjected to fluorescent imaging before and after staining with Sybr gold.

In case of 5S rRNA labeling using FJC9 ribozyme, 50 ng of the total cellular RNA was mixed with 100 pmol of the corresponding FJC9 ribozyme. Reaction was done in 5 μ L of the selection buffer including 40 mM MgCl₂ and 200 μ M, *N*⁶-Fluorescein-TenDP. The reaction was incubated at 37°C for 6 h. Afterwards, the reaction was subjected to ethanol precipitation and the resulting pellet was dissolved in 2 μ L of milliQ water and 2 μ L of the stop solution. The sample was resolved on a 10% analytical denaturing gel, next to the fluorescein labeled in vitro transcribed 5S rRNA as size marker. The gel was then subjected to fluorescent imaging before and after staining with Sybr gold.

2.3.4 Mutually orthogonal dual-color labeling of a synthetic transcript

A synthetic substrate was designed by tandem fusion of the parent substrate sequence and one of its derivatives via a 4 nt linker. The 41 nt RNA was prepared by in vitro transcription with T7 RNA polymerase. 20 pmol of this substrate sequence was mixed with 50 pmol of each ribozyme in a total volume of 5 μ L of selection buffer containing 40 mM MgCl₂ and 200 μ M of either Cy5-TenDP or 6-FAM-ATP or both. The FJ1 ribozyme was targeted towards the parent substrate sequence while the FH14 was designed to target the mutated segment. Time-point samples (0.5 μ L) were taken at 0, 5 h and after overnight incubation and quenched by adding to 99.5 μ L of TEN buffer. The samples were then subjected to ethanol precipitation, and the pellet was dissolved in 5 μ L of the stop solution. 2.5 μ L of these samples were then resolved on 15% analytical denaturing PAGE for 45 min and subjected to dual-channel fluorescent imaging. To confirm the specificity of each ribozyme for its cognate NTP analog and its target sequence an 8-17NG deoxyribozyme was designed to cleave the double-labeling RNA substrate asymmetrically resulting in a 24 nt fragment expected to harbor the label installed by FJ1 (Cy5-tenofovir branch) and a 17 nt fragment supposed to carry the label attached by FH14 (6-FAM-AMP branch). To perform the cleavage experiment, an aliquot (3.5 μ L) of the labeling reactions explained above was subjected to ethanol precipitation. The pellet was then dissolved in 10 μ L of 8-17NG reaction buffer containing 50 mM HEPES (pH= 7.5), 400 mM KCl, 100 mM NaCl, 10 mM MnCl₂, 10 mM MgCl₂ and 200 pmol of the corresponding 8-17NG deoxyribozyme. The reaction was then incubated at 37°C and 1.4 μ L timepoint samples were taken at 0, 0.5 and 1 h and mixed with 3.57 μ L of the stop solution. 2.5 μ L of these samples were resolved on the same gel as the double labeling reaction.

SUPPORTING INFORMATION

2.3.5. Probing of the modification site on the target cellular RNA using reverse transcription**2.3.5.1. Modification of the cellular RNA for primer extension experiments**

200 ng of *E. coli* total cellular RNA and 20 pmol of individual ribozymes were dissolved in a total volume of 10 μ L of the selection buffer including 40 mM $MgCl_2$ and the ribozyme's cognate substrate. For FH14 type ribozyme N^6 -aminoethyl-ATP- biotin (200 μ M final concentration) was used, and for the FJ1 type ribozyme Biotin-Tenofovir-DP-biotin (7, 340 μ M final concentration). The reactions were incubated at 37°C for 5 h and were subjected to ethanol precipitation. The pellet was dried and directly used in the primer extension reaction.

2.3.5.2. ^{32}P -Labeling of the primers

Primers (100 pmol) were dissolved in 10 μ L of PNK buffer A including 5 μ Ci of γ - ^{32}P -ATP, and 5 units of PNK. The reaction was incubated at 37°C for 1 h, followed by two rounds of ethanol precipitation. The pellet was dissolved in 10 μ L of milliQ water, resulting in a sample with 2000-3500 IPS.

2.3.5.3. Primer extension protocol

5 pmol of the ^{32}P -labeled primer and 200 ng of the ribozyme-modified or non-modified total cellular RNA were dissolved in 5 μ L of the annealing buffer (Tris 5 mM pH= 7.5, EDTA 0.1 mM). The sample was then placed at 95°C for 3 minutes, followed by incubation at room temperature for 10 minutes. 2 μ L of the 5X First strand synthesis buffer (Invitrogen), 0.5 μ L of 0.1 M DTT, 0.5 μ L of 10 mM dNTP mix and 50 units of superscript III reverse transcriptase was then added to the reaction and the final volume was then adjusted to 10 μ L by adding milliQ water. The reaction was then incubated at 55°C for 1 hour. 1 μ L of 2 N NaOH was then added to the reaction and the sample was incubated for 5 minutes at 95°C. The reaction was then subjected to ethanol precipitation. The dried pellet was then dissolved in 10 μ L of high-dye loading buffer. 2.5 μ L of this sample was then resolved using 15%, analytical 45 cm long PAGE, under constant power of 45 W for 2 hours and 15 minutes. The gel was dried at 80°C under vacuum for 30 minutes and exposed to the phosphorus screen overnight. The screen was then scanned on a Typhoon phosphorus imager.

The sequencing reactions were performed by annealing 5 pmol of the ^{32}P -labeled primers to 200 ng of the non-modified total cellular RNA in 5 μ L of the annealing buffer. for the A sequencing ladder ddTTP, was added to the reaction to a final concentration of 0.5 mM and dTTP at 0.05 mM. The rest of the dNTP were adjusted to a final concentration of 0.5 mM. For G sequencing ladder ddCTP was included in the reaction at a concentration of 0.5 mM and dCTP at 0.05 mM. The rest of the dNTPs had a final concentration of 0.5 mM. All the other reagents such as the buffer conditions and enzyme units were the same as the primer extension protocol.

SUPPORTING INFORMATION

3. Results and Discussion

3.1 Sequencing results of FJ ribozyme selection

3.1.1. FJ ribozyme sequences identified by Sanger sequencing.

Forty colonies were checked by colony-PCR, 10 plasmids were isolated and sequenced

FJ1 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC
 FJ2 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC
 FJ3 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC
 FJ4 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC
 FJ24 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC
 FJ32 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC
 FJ35 TTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTATGTCC

FJ8 YTGAAGGC ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC CAGTATGTCC
 FJ11 TTGAAGGC ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC CAGTATGTCC

3.1.2. FJ ribozyme sequences identified by Illumina sequencing.

After quality control and trimming of illumina adapters the sequencing data were analyzed by fastaptamer.^[9]

fastaptamer_count: FJ_Round 7: 4.789.062 total reads, 288.126 unique sequences
 FJ_Round 12: 1.138.422 total reads, 66.443 unique sequences

fastaptamer_cluster: FJ_Round 12

Cluster	Unique	#Reads	RPM
1	1407	866354	761012.5
2	407	84781	74472.2
3	259	55156	48449.4
4	158	21523	18905.9
5	28	3697	3247.4
6	38	3943	3463.5
7	20	2733	2400.6
8	26	2401	2109.0
9	10	1268	1113.8
10	6	720	632.4

fastaptamer_compare: RPM(x) = Round 7 and RPM(y) = Round 12

First sequence of first ten clusters ordered by log₂ (enrichment)

Cluster	sequence	RPM(x)	RPM(y)	log ₂ x/y
>2-2-1-0=FJ8	AACCAGCCTACCATCCTTGAAGGC ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC CAGT	38.8	33325.9	9.74
>8-4-1-0=FJ1	AACCAGCCTACCATCCTTGAAGGC CCACCCTCATAAACTGAAGATCCTTTGGCAAGGGTCTA CAGTA	25.7	10479.4	8.67
>82-9-1-0=FJC9	AACCAGCCTACCATCCTTGAAGGC ACGAGATATGTTGCACTACTTTAGCGAATTGGGCATCC CAGT	46.2	924.1	4.32
>1-1-1-0=FJC1	AACCAGCCTACCATCCTTGAAGGC GAAACGTGTACATAAGAAACGGTAAACTAGCAAGTTC CAGT	130185	502593	1.95
>48-7-1-0 (a)	AACCAGCCTACCATCCTTGAAGGC T CAGTATGTC GACCTTCTGTCTTATACACATCTCCGAGCCCA	474.2	1586.4	1.74
>26-5-1-0 (b)	AACCAGCCTACCATCCTTGAAGGC CGAATGCCACCGAA CAGTATGTC GACCTTCTGTCTTATACAC	3096.4	2437.6	-0.35
>3-3-1-0=FJC3	AACCAGCCTACCATCCTTGAAGGC CGAATGCCACCGAACCCTGATAATTGGCGCTCCAATTTC CAGTA	39964.6	30094.3	-0.41
>56-8-1-0	AACCAGCCTACCATCCTTGAAGGC ACTGGTGTACAAAGTAAACGCAACCTTAATTTTACGATCC CAGT	4264.5	1258.8	-1.76
>32-6-1-0 (c)	AACCAGCCTACCATCCTTGAAGGC ATCGGTGTAACATAAACGAAACCTTAGGTCCTTTGGCCCC CAGT	34139.0	2206.6	-3.95
>126-10-1-0	AACCAGCCTACCATCCTTGAAGGC ATAGTTTCAACAATAGCCACTTAAGCTTATCCAAAGCTCA CAGT	11488.7	577.1	-4.32

(a) = extended complement to substrate sequence

(b) reveals a fully complementary binding arm after 14 nt in the center of N40 core

(c) related to FJ8

SUPPORTING INFORMATION

FJ8 (407 sequences)	YTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGTATGTCC
>2-37939-33325.95-2-1-0	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>5-18208-15994.07-2-2-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CAGT
>52-1677-1473.09-2-3-2	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CA
>91-893-784.42-2-4-2	ACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGTA
>97-855-751.04-2-5-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>101-840-737.86-2-6-3	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CA
>108-800-702.73-2-7-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CAGT
>122-673-591.17-2-8-3	AACCAGGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CA
>141-519-455.89-2-9-4	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACCTTACCTTCCCC	CAGTAT
>153-468-411.10-2-10-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>161-417-366.30-2-11-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>162-415-364.54-2-12-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>167-404-354.88-2-13-3	ACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CAGTA
>170-387-339.94-2-14-2	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CAGT
>184-341-299.54-2-15-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>198-300-263.52-2-16-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>200-286-251.22-2-17-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>202-285-250.35-2-18-4	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CA
>214-255-223.99-2-19-1	AACCAGCCTACCATCCTTGAAGGC	ATGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>217-252-221.36-2-20-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACGAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>219-247-216.97-2-21-2	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CAGT
>223-239-209.94-2-22-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
>227-235-206.43-2-23-2	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	AGTA
>228-232-203.79-2-24-1	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGT
...			
>2192-11-9.66-2-405-2	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACAGTACCTTCCCC	CAGT
>2192-11-9.66-2-406-3	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CA
>2192-11-9.66-2-407-2	AACCAGCCTACCATCCTTGAAGGC	ACGGTGTTACAAAACGAAACCTTAGGACATTACCTTCCCC	CAGTA
FJ1 (158 sequences)	TTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTATGTCC
>8-11930-10479.42-4-1-0	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>63-1338-1175.31-4-2-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGT
>64-1316-1155.99-4-3-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>128-653-573.60-4-4-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>150-471-413.73-4-5-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>191-319-280.21-4-6-2	ACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTAT
>216-253-222.24-4-7-2	AACCAGGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGT
>230-228-200.28-4-8-2	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTACAGT	
>283-165-144.94-4-9-2	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGT
>297-158-138.79-4-10-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTTATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>302-156-137.03-4-11-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>347-127-111.56-4-12-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>355-121-106.29-4-13-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>417-96-84.33-4-14-2	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGT
>417-96-84.33-4-15-1	AACCAGCCTACCATCCTTGAAGGC	CCATCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>427-93-81.69-4-16-2	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTAT
>433-92-80.81-4-17-1	AACCAGCCTACCATCCTTGAAGGC	CCACTCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>436-91-79.94-4-18-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTA
>502-75-65.88-4-19-3	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CA
>502-75-65.88-4-20-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAGATCCTTTGGCAAGGGTCTA	CAGTA
...			
>2192-11-9.66-4-156-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCATTGGCAAGGGTCTA	CAGTA
>2192-11-9.66-4-157-2	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGT
>2192-11-9.66-4-158-1	AACCAGCCTACCATCCTTGAAGGC	CCACCCTCATAAAACTGAAGATCCTTTGGCAAGGGTCTA	CAGTC
FJC9 (10 sequences)			
>82-1052-924.09-9-1-0	AACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGT
>701-50-43.92-9-2-1	AACCAGCCTACCATCCTTGAAGGC	ACAAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGT
>918-33-28.99-9-3-2	AACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CA
>981-30-26.35-9-4-1	AACCAGCCTACCATCCTTGAAGGC	ACTAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGT
>1146-24-21.08-9-5-3	AACCAGGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CA
>1391-19-16.69-9-6-1	AACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCA	CAGT
>1457-18-15.81-9-7-2	ACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGTA
>1520-17-14.93-9-8-1	AACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGT
>1927-13-11.42-9-9-1	AACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGT
>2058-12-10.54-9-10-1	AACCAGCCTACCATCCTTGAAGGC	ACGAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC	CAGT

Partial alignment of N40 core:

FJC1 GAAACGTG-TCACATAA--GAAAA-CGGTAAACTAGCAAGTTCC
 FJC3 CGAA---TGCCACCG-AACCGTATAA-TTGCCGCCCTCCAATTC
 FJC9 ACAGATATGTTGCACTACACTTTAGCGAATTGGGCATCC

FJ8 A-CGGTGTTACA---AAACGAAACCTTAGGACATTACCTTCCCC
 FJC6 ATCGGTGTAACAT--AAACGAAACCTTAGG--TCCTTGGCCCC
 FJC8 ACTGGTGTTACAAGTAAACGCAACCTTA---ATTTTACGATCC

SUPPORTING INFORMATION

3.2 Supporting Figures




Figure S1. Biotin-Tenofovir diphosphate is not a substrate for FH14. (3'-fluorescein-labeled parent substrate RNA; 200 μ M N^6 -Biotin-ATP, 800 μ M N^6 -Biotin-TenDP, with 40 mM $MgCl_2$, in selection buffer. Overnight incubation at 37°C.




Figure S2. In vitro selection scheme, showing RNA substrate sequence in red, random region (N_{40}) in green, connecting loop in cyan, and in purple are the ribozyme binding arms complementary to the substrate sequence (flanking the bulged A). The primer binding sites for the first PCR are indicated. A second PCR generated the dsDNA template containing the T7 promoter (light green) necessary for transcription of the RNA.

SUPPORTING INFORMATION




Figure S3. Expanded version of Figure 2b with legend of color-coded clusters from fastaptamer_cluster FJ_round12. See also Section 3.1.2

SUPPORTING INFORMATION




Figure S4. Secondary structures and dotplots of FJ1, FJ8 and FJC9. Dot plots were generated by Vienna RNA package.^[10] The minimum free energy secondary structure is drawn; for FJ1 and FJ8 one possible alternative secondary structure is also shown.

SUPPORTING INFORMATION




Figure S5. ESI mass spectrometric analysis of biotin-tenofovir-labeled 3'-aminohexyl-RNA produced by preparative labelling with FJ1. ESI-Mass spectrum (neg. mode) of the isolated RNA product; deconvoluted MS (red) and simulated mass spectrum (grey).

SUPPORTING INFORMATION




Figure S6. Alkaline hydrolysis and RNase T1 digestion of isolated FJ1 product for assignment of labeling site. Experimental details see section 2.3.2.2.




Figure S7. Gel images for data in Figure 4a. Small molecule reactivity with FJ1, FJ8 and FJC9 on 3'-fluorescein-labeled parent RNA. (300 μ M Ten-DP analog, with 40 mM $MgCl_2$, in selection buffer, timepoints are: 0, 15, 30, 60 min; 2, 4, 6, 8, 22 h).

SUPPORTING INFORMATION




Figure S8. Substrate sequence mutations for FJC9. (3'-fluorescein-labeled RNAs, 300 μ M Biotin-Ten-DP, with 40 mM MgCl₂, in selection buffer, 37°C, timepoints are: 0, 15, 30, 60 min; 2, 4, 6, 8, 22 h).




Figure S9. Substrate sequence scope for FJ1. Transition and transversion mutations are color coded in the sequence labels: pink: transition, purple: transversion-1, blue: transversion-2. Green: nucleotides that remained unchanged from the parent. 5'-GG in grey, since these are needed for transcription initiation. (3'-fluorescein-labeled RNAs, 300 μ M Biotin-Ten-DP, with 40 mM MgCl₂, in selection buffer, 37°C, timepoints are: 0, 15, 30, 60 min; 2, 4, 6, 8, 22 h).

SUPPORTING INFORMATION




Figure S10. Mismatch effects on ribozyme labeling efficiency. The substrates were single point mutants with the central GAG motif changed to AAG or GAA. Labeling reactions were performed with parent FJ1 and FJC9 ribozymes, thus introducing 5' or 3' A-C mismatches. In both cases, the labeling efficiency was significantly reduced (below 15% after 22 h, compared to 90% in the matched case). This result indicates that there is little propensity for off-target labeling, i.e. both ribozymes require full complementarity to maintain efficient labeling.




Figure S11. a) Orthogonality of FJC9 and FJ1 to FH14. FJ ribozymes do not show any RNA labelling activity with ATP. (3'-fluorescein-labeled parent RNA, 300 μ M Biotin-Ten-DP, 200 μ M Biotin-ATP, with 40 mM $MgCl_2$, in selection buffer, 37°C, timepoints are: 0, 1, 2, 4, 6, 22 h). b) Mg^{2+} -dependence of FJ1 ribozyme. Reactions were performed with parent RNA and biotin-tenofovir-DP at 37°C for 1, 6, and 22 h. The ribozyme activity is dependent on Mg^{2+} concentration. At 5 mM Mg^{2+} , only 7 % labeled RNA is observed after 22h, but already at 10 mM Mg^{2+} , the yield is above 60%.

SUPPORTING INFORMATION




Figure S13. Labelling of *E. coli* 16S and 23S rRNA with FJ1 and FH14. Lane 1: unlabelled reference. lanes 2-4: FJ1 with Cy5-Ten-DP on three different positions of 16S rRNA, lanes 5-7: FH14 with Atto550-ATP on 3 different positions of 23S rRNA, lane 8: simultaneous labelling of 16S and 23S rRNA. lanes 9-11: FJ1 with Cy5-Ten-DP on three different positions of 23S rRNA, lane 12-13: FH14 with FAM-ATP on two different positions of 16S rRNA.




Figure S14. Primer extension analysis of FJ1 and FH14 labelling sites in 16S and 23S rRNA. As indicated on top of each gel picture, *E. coli* RNA was labelled by FJ1 and FH14 ribozymes targeting positions A649, A383 and A325 in 16S rRNA and A637 in 23S rRNA, respectively, using cognate biotinylated tenofovir or ATP substrates. Primer extension assays on the modified RNA and the unmodified input RNA were performed with $5'$ - ^{32}P -labelled primers and superscript III reverse transcriptase. The primer sequences are listed in section 1.2. Sequencing lanes (A, G) are indicated.




Figure S15. Comparison of *R*-enantiomer and racemic tenofovir diphosphate for RNA labelling reaction with FJ1 on parent RNA substrate. (last timepoint is 22h).

SUPPORTING INFORMATION

HPLC chromatograms of FAM-Ten-DP (17)






Figure S16: HPLC chromatograms of compound 13, FAM azide and product 17 at 260 nm (above) and at 496 nm (below), respectively.

SUPPORTING INFORMATION

HPLC chromatograms of Cy5-Ten-DP (18)









Figure S17: HPLC chromatograms of compound **13**, SulfoCy5 azide and product **18** at 260 nm (above) and at 647 nm (below), respectively.

SUPPORTING INFORMATION



NMR spectra

Compound 1 ^1H NMRCompound 1 ^{13}C NMR



SUPPORTING INFORMATION

Compound 2 ¹H NMRCompound 2 ¹³C NMR



SUPPORTING INFORMATION

Compound 3 ^1H NMRCompound 3 ^{13}C NMR



SUPPORTING INFORMATION

Compound 3 ^{31}P NMRCompound 4 ^1H NMR



SUPPORTING INFORMATION

Compound 4 ^{13}C NMRCompound 4 ^{31}P NMR



SUPPORTING INFORMATION

Compound 5 ^1H NMRCompound 5 ^{13}C NMR


SUPPORTING INFORMATION

Compound 5 ^{31}P NMRCompound 6 ^1H NMR



SUPPORTING INFORMATION

Compound 6 ^{31}P NMRCompound 7 ^1H NMR



SUPPORTING INFORMATION

Compound 7 ^{31}P NMR



SUPPORTING INFORMATION

Compound 8 ^1H NMRCompound 8 ^{13}C NMR



SUPPORTING INFORMATION

Compound **8** ^{31}P NMRCompound **8b** ^1H NMR


SUPPORTING INFORMATION

Compound 8b ^{13}C NMRCompound 8b ^{31}P NMR



SUPPORTING INFORMATION

Compound 9 ^1H NMRCompound 9 ^{13}C NMR



SUPPORTING INFORMATION

Compound 10 ^{31}P NMR



SUPPORTING INFORMATION

Compound 11 ¹H NMRCompound 11 ¹³C NMR



SUPPORTING INFORMATION

Compound 11 ^{31}P NMRCompound 12 ^1H NMR



SUPPORTING INFORMATION

Compound 12 ^{13}C NMRCompound 12 ^{31}P NMR



SUPPORTING INFORMATION

Compound 13 ^1H NMRCompound 13 ^{31}P NMR



SUPPORTING INFORMATION

Compound 14 ¹H NMRCompound 14 ¹³C NMR



SUPPORTING INFORMATION

Compound 14 ^{31}P NMR:Compound 15 ^1H NMR:



SUPPORTING INFORMATION

Compound 15 ^{13}C NMRCompound 15 ^{31}P NMR



SUPPORTING INFORMATION

Compound 19 ¹H NMRCompound 19 ¹³C NMR

SUPPORTING INFORMATION

Compound 20 ^1H NMRCompound 20 ^{13}C NMR

SUPPORTING INFORMATION

Compound 21 ^1H NMRCompound 21 ^{13}C NMR

References

- [1] M. Ghaem Maghami, C. P. M. Scheitt, C. Höbartner, *J Am Chem Soc* **2019**, *141*, 19546-19549.
- [2] J. C. Kim, S. H. Kim, J. A. Kim, S. K. Choi, W. W. Park, *Arch Pharm Res* **1998**, *21*, 458-464.
- [3] A. L. Jeffery, J. H. Kim, D. F. Wiemer, *Tetrahedron* **2000**, *56*, 5077-5083.
- [4] Z. Liu, N. Yasuda, M. Simeone, R. A. Reamer, *J Org Chem* **2014**, *79*, 11792-11796.
- [5] a) D. E. Hoard, D. G. Ott, *J Am Chem Soc* **1965**, *87*, 1785-1788; b) E. S. Simon, S. Grabowski, G. M. Whitesides, *Journal of Organic Chemistry* **1990**, *55*, 1834-1841.
- [6] L. M. Schultze, H. H. Chapman, N. J. P. Dubree, R. J. Jones, K. M. Kent, T. T. Lee, M. S. Louie, M. J. Postich, E. J. Prisbe, J. C. Rohloff, R. H. Yu, *Tetrahedron Lett* **1998**, *39*, 1853-1856.
- [7] C. M. Leenders, G. Jansen, M. M. Frissen, R. P. Lafleur, I. K. Voets, A. R. Palmans, E. W. Meijer, *Chemistry* **2016**, *22*, 4608-4615.
- [8] M. B. Stead, A. Agrawal, K. E. Bowden, R. Nasir, B. K. Mohanty, R. B. Meagher, S. R. Kushner, *Nucleic Acids Res* **2012**, *40*, e156.
- [9] K. K. Alam, J. L. Chang, D. H. Burke, *Mol Ther Nucleic Acids* **2015**, *4*, e230.
- [10] <http://ma.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi>.
- [11] 16S rRNA: <https://macentral.org/rna/URS000004ABFE9/562>; 23S rRNA: <https://macentral.org/rna/URS00004B0F34/562>;