

Impact of the Preparation Procedure on the Performance of the Microporous HKUST-1 Metal-Organic Framework in the Liquid-Phase Separation of Aromatic Compounds

Vera I. Isaeva ^{1,2,*}, Bulat R. Saifutdinov ^{3,*}, Vladimir V. Chernyshev ^{3,4}, Vadim V. Vergun ¹, Gennady I. Kapustin ^{1,2}, Yulia P. Kurnysheva ³, Mikhail M. Ilyin ⁵ and Leonid M. Kustov ^{1,2}

¹ N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, Leninsky prospect 47, Moscow 119991, Russia; polubrat@mail.ru (V.V.V.); gik@ioc.ac.ru (G.I.K.); lmkustov@mail.ru (L.M.K.)

² National University of Science and Technology MISiS, Leninsky prospect 4, Moscow 119991, Russia

³ A.N. Frumkin Institute of Physical Chemistry and Electrochemistry, Russian Academy of Sciences, Building 4, 31 Leninsky prospect, Moscow 119071, Russia; vladimir@struct.chem.msu.ru (V.V.C.); kurnyshevay@mail.ru (Y.P.K.)

⁴ Department of Chemistry, M.V. Lomonosov Moscow State University, 1-3 Leninskie Gory, Moscow 119991, Russia

⁵ A.N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 28 Vavilov St., Moscow 119991, Russia; mil@ineos.ac.ru (M.M.I.)

* Correspondence: sharf@ioc.ac.ru (V.I.I.); bulatrsaifutdinov@gmail.com (B.R.S.)




Figure S1. Output curve recorded during the elution of 2,3-dichlorophenol at 65°C through the porous HKUST-1 layer.




Figure S2. Output curve recorded during the elution of 2,4-dichlorophenol at 65°C through the porous HKUST-1 layer.




Figure S3. Output curve recorded during the elution of 2,5-dichlorophenol at 65°C through the porous HKUST-1 layer.




Figure S4. Output curve recorded during the elution of 2,6-dichlorophenol at 65°C through the porous HKUST-1 layer.