

Table S3a. Factors associated with correct and incorrect use of mCIDT for each diagnosis, categorical variables (n=34)

	Psychosis			Alcohol Use Disorder			Postpartum Depression			Epilepsy		
	Incorrect mCIDT, n (%)	Correct mCIDT, n (%)	p-value	Incorrect mCIDT, n (%)	Correct mCIDT, n (%)	p-value	Incorrect mCIDT, n (%)	Correct mCIDT, n (%)	p-value	Incorrect mCIDT, n (%)	Correct mCIDT, n (%)	p-value
Education			.000			.000			.000			.000
Illiterate	1(0)	3(100)		1(25)	3(75)		2(50)	2(50)		1(25)	3(75)	
Literate	17(89.5)	2(10.5)		17(89)	2(10)		17(89)	2(11)		16(84.2)	3(15.3)	
Grade 1-5	2(66.7)	1(33.3)		0(0)	3(100)		1(33.3)	2(66.7)		0(0)	3(100)	
Grade 5-10	0(0)	3(100)		0(0)	3(100)		1(33.3)	2(66.7)		0(0)	3(100)	
SLC pass & above	0(0)	5(100)		0(0)	5(100)		0(0)	5(100)		0(0)	5(100)	
Ward			.267			.368			.184			.168
Ratanchura (1)	4(44.4)	5(55.6)		4(44.4)	5(55.6)		4(44.4)	5(55.6)		2(22.2)	7(77.8)	
Bhimeshwor (2)	4(44.4)	5(55.6)		3(33)	6(66)		4(44.4)	5(55.6)		4(44.4)	5(55.6)	
Bhuwaneshwori (3)	7(87.5)	1(12.5)		6(75)	2(25)		6(75)	2(25)		5(62.5)	3(37.5)	
Basehwor (4)	5(62.5)	3(37.5)		5(62.5)	3(37.5)		7(87.5)	1(12.5)		6(75)	2(25)	
Can send a SMS			.017			.003			.039			.015
No	13(81.2)	3(18.7)		13(81)	3(19)		13(81)	3(19)		12(75)	4(25)	
Yes	7(38)	11(62)		5(27)	13(73)		8(44.4)	10(55.6)		5(27.8)	13(72.2)	
Can use mCIDT Codebook			.004			.000			.012			.084
No	15(83.3)	3(16.7)		15(83)	3(17)		15(83)	3(17)		12(66.7)	6(33.3)	
Yes	5(31.3)	11(68.7)		3(18.8)	13(81.3)		6(37.5)	10(62.5)		5(31.3)	11(68.8)	

Two-tailed Fisher's exact test t-test compares correctly or incorrectly completing mCIDT step 1 message completion among persons who accurately identified paper CIDT.

Table S3b. Factors associated with correct and incorrect use of mCIDT for each diagnosis, continuous variables (n=34)

	Psychosis				Alcohol Use Disorder				Postpartum Depression				Epilepsy			
	Incorrect mCIDT, Median (IQR)	Correct mCIDT, Median (IQR)	Test statistic (Wilcoxon)	p-value	Incorrect mCIDT, Median (IQR)	Correct mCIDT, Median (IQR)	Test statistic (Wilcoxon)	p-value	Incorrect mCIDT, Median (IQR)	Correct mCIDT, Median (IQR)	Test statistic (Wilcoxon)	p-value	Incorrect mCIDT, Median (IQR)	Correct mCIDT, Median (IQR)	Test statistic (Wilcoxon)	p-value
Age (years)	57 (48, 60)	40 (35, 51.25)	17.50	.009	52.5 (47.25, 59.5)	42 (35, 48.75)	40.00	.001	51.5 (44.8, 60)	45 (36, 48.5)	57.00	.016	53 (46.5, 60.5)	45 (36, 52)	67.00	.08