

Supplemental Table 2a: Multivariate Regression Model of Variables Predicting 30-Day Continence

	B	S.E.	Wald	Sig.	OR	95% CI	
						Lower	Upper
MUL (≤ 1.4 cm [ref] v. > 1.4 cm)	1.579	0.695	5.157	0.023	4.851	1.241	18.961
Age (cont.)	-0.078	0.030	6.910	0.009	0.925	0.873	0.980
pGS (< 8 v. 8-10 [ref])	0.261	0.559	0.219	0.640	1.299	0.434	3.883
pT2, positive margins	0.200	0.437	0.209	0.647	1.221	0.519	2.873
Pre-op PSA (< 10 v. > 10 [ref])	-0.021	0.026	0.655	0.418	0.979	0.930	1.031
Pre-op IIEF-5 (cont.)	0.021	0.034	0.399	0.528	1.021	0.956	1.091
BMI (cont.)	0.047	0.073	0.422	0.516	1.048	0.909	1.209
Constant	1.788	3.141	0.324	0.569	5.976		

Supplemental Table 2b: Multivariate Regression Model of Variables Predicting 1-Year Continence

	B	S.E.	Wald	Sig.	OR	95% CI	
						Lower	Upper
MUL (≤ 1.4 cm [ref] v. > 1.4 cm)	2.421	1.201	4.065	0.044	11.255	1.070	118.407
Age (cont.)	-0.085	0.072	1.368	0.242	0.919	0.797	1.059
pGS (< 8 v. 8-10 [ref])	-0.354	1.211	0.086	0.770	0.702	0.065	7.527
pT2, positive margins	-0.534	0.971	0.303	0.582	0.586	0.087	3.934
Pre-op PSA (< 10 v. > 10 [ref])	0.019	0.074	0.063	0.801	1.019	0.881	1.179
Pre-op IIEF-5 (cont.)	0.099	0.069	2.059	0.151	1.104	0.964	1.264
BMI (cont.)	0.089	0.189	0.223	0.637	1.093	0.755	1.584
Constant	3.639	7.502	0.235	0.628	38.046		