





Rapid onset honeycombing fibrosis in spontaneously breathing patient with COVID-19

Margot Combet¹, Arthur Pavot ¹, Laurent Savale ^{2,3,4}, Marc Humbert ^{2,3,4} and Xavier Monnet^{1,2,4}

Affiliations: ¹AP-HP, Service de médecine intensive-réanimation, Hôpital de Bicêtre, Le Kremlin-Bicêtre, France. ²Université Paris-Saclay, Faculté de Médecine, Le Kremlin-Bicêtre, France. ³AP-HP, Service de pneumologie et soins intensifs respiratoires, Centre de Référence de l'Hypertension Pulmonaire, Hôpital de Bicêtre, Le Kremlin-Bicêtre, France. ⁴INSERM UMR_S 999, Hôpital Marie-Lannelongue, Le Plessis-Robinson, France.

Correspondence: Arthur Pavot, AP-HP, Service de médecine intensive-réanimation, Hôpital de Bicêtre, 63 rue Gabriel Péri, Le Kremlin-Bicetre, 94275, France. E-mail: pavot.arthur@gmail.com

@ERSpublications

Patients with COVID-19 should be monitored for rapid onset honeycombing fibrosis https://bit.ly/2B7fRcA

Cite this article as: Combet M, Pavot A, Savale L, *et al.* Rapid onset honeycombing fibrosis in spontaneously breathing patient with COVID-19. *Eur Respir J* 2020; 56: 2001808 [https://doi.org/10.1183/13993003.01808-2020].

This single-page version can be shared freely online.

To the Editor:

A 38-year-old man presented with a 6-day history of gradually worsening cough and asthenia and secondary anosmia and agueusia. He had no prior personal medical history. His mother had a history of diabetes mellitus and his son was followed for haemophilia A. His spouse had exhibited similar symptoms a week earlier.

Copyright ©ERS 2020.. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.