

Supplementary Online Content

Jabri A, Kalra A, Kumar A, et al. Incidence of stress cardiomyopathy during the coronavirus disease 2019 pandemic. *JAMA Netw Open*. 2020;3(7):e2014780. doi:10.1001/jamanetworkopen.2020.14780

eTable. Incidence Rate Ratios of Stress Cardiomyopathy Comparing the COVID-19 Pandemic Period With Prepandemic Periods

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Incidence Rate Ratios of Stress Cardiomyopathy Comparing the COVID-19 Pandemic Period With Prepandemic Periods

| | Risk Ratio | 95% CI | P value |
|---|-------------------|---------------|----------------|
| Unadjusted Models | | | |
| COVID19 vs. all pre-COVID19 | 4.58 | (4.11, 5.11) | <.0001 |
| COVID19 (Mar - Apr 2020) vs. Jan - Feb 2020 | 4.31 | (1.62, 11.48) | 0.004 |
| COVID19 (Mar - Apr 2020) vs. Mar - Apr 2019 | 4.39 | (2.14, 8.97) | <.0001 |
| COVID19 (Mar - Apr 2020) vs. Jan - Feb 2019 | 4.79 | (1.80, 12.77) | 0.002 |
| COVID19 (Mar - Apr 2020) vs. Mar - Apr 2018 | 5.04 | (2.02, 12.55) | 0.0005 |
| Adjusted Models | | | |
| HTN: COVID19 vs. all pre-COVID19 | 5.01 | (4.55, 5.51) | <.0001 |
| HLD: COVID19 vs. all pre-COVID19 | 4.68 | (4.05, 5.40) | <.0001 |
| CAD: COVID19 vs. all pre-COVID19 | 4.79 | (4.31, 5.32) | <.0001 |
| Asthma/COPD: COVID19 vs. all pre-COVID19 | 4.16 | (3.87, 4.47) | <.0001 |
| CAD = coronary artery disease, CI = confidence interval, COPD = chronic obstructive pulmonary disease, HLD = hyperlipidemia, HTN = hypertension | | | |