

Article

Retrospective Multicentric Study on Non-Optic CNS Tumors in Children and Adolescents with Neurofibromatosis Type 1

Claudia Santoro, Stefania Picariello, Federica Palladino, Pietro Spennato, Daniela Melis, Jonathan Roth, Mario Cirillo, Lucia Quaglietta, Alessandra D'Amico, Giuseppina Gaudino, Maria Chiara Meucci, Ursula Ferrara, Shlomi Constantini, Silverio Perrotta and Giuseppe Cinalli

Supplementary Table S1. Univariate Cox regression analysis of progression free survival in patients with low grade gliomas.

	Hazard Ratio (95%CI)	<i>p</i>
Multiple lesions	2.81 (1.06–7.46)	0.038
Optic pathway glioma co-presence	1.24 (0.52–2.97)	0.629
Surgery first approach	0.48 (0.22–1.04)	0.062
Age at diagnosis (continuous variable)	0.99 (0.99–1.01)	0.902
Age at diagnosis < 10 years	1.12 (0.46–2.71)	0.803
Female gender	1.27 (0.81–2.00)	0.292
Inherited NF1	1.37 (0.86–2.19)	0.189
Symptomatic at diagnosis	0.46 (0.17–1.26)	0.131
Brainstem vs other locations	0.73 (0.28–1.90)	0.519
Posterior fossa vs other locations	1.07 (0.38–3.03)	0.895