

Sonication-assisted production of fosetyl-Al nanocrystals: investigation of human toxicity and *in vitro* antibacterial efficacy against *Xylella fastidiosa*

Supplementary Materials

Francesca Baldassarre ^{1,2,*†}, Giuseppe Tatulli ^{3†}, Viviana Vergaro ^{1,2}, Stefania Mariano ⁴, Valeria Scala ³, Concetta Nobile ², Nicoletta Pucci ³, Luciana Dini ^{2,5}, Stefania Loreti ³ and Giuseppe Ciccarella ^{1,2,*}

¹ Biological and Environmental Sciences Department, Udr INSTM of Lecce University of Salento, Via Monteroni, 73100 Lecce, Italy; viviana.vergaro@unisalento.it

² Institute of Nanotechnology, CNR NANOTEC, Consiglio Nazionale delle Ricerche, Via Monteroni, 73100 Lecce, Italy; concetta.nobile@nanotec.cnr.it (C.N.); luciana.dini@uniroma1.it (L.D.);

³ Council for Agricultural Research and Economics, Research Centre for Plant Protection and Certification of Rome, 00156 Rome, Italy; giuseppe.tatulli@hotmail.it (G.T.); valeria.scala@crea.gov.it (V.S.); nicoletta.pucci@crea.gov.it (N.P.); stefania.loreti@crea.gov.it (S.L.)

⁴ Biological and Environmental Sciences Department, University of Salento, Via Monteroni, 73100 Lecce, Italy; stefania.mariano@unisalento.it

⁵ Department of Biology and Biotechnology “Charles Darwin”, University of Rome “La Sapienza”, Piazzale Aldo Moro 5, 00185 Roma, Italy

* Correspondence: francesca.baldassarre@unisalento.it (F.B.); giuseppe.ciccarella@unisalento.it (G.C.); Tel.: +39-0832-319207 (F.B.); +39-0832-319810 (G.C.)

† These authors contributed equally to this work.




Figure S1. Highly magnified detail of a wrinkled structure in the SEM image of figure 3D.




Figure 2. TEM image of a CH-nanoFos colloid.