

Supplementary Materials

Novel Two Terpenes Isolated from *Dictyostelium*

Cellular Sime Molds

Hitomi Sasaki,¹ Yuzuru Kubohara,² Hirotaka Ishigaki,³ Katsunori Takahashi,³ Hiromi Eguchi,¹

Akihiro Sugawara,¹ Yoshiteru Oshima¹ and Haruhisa Kikuchi^{1,*}

¹ Graduate School of Pharmaceutical Sciences, Tohoku University, 6-3, Aza-Aoba, Aramaki, Aoba-ku, Sendai 980-8578, Japan.

² Graduate School of Health and Sports Science, Juntendo University, 1-1 Hiraga-gakuendai, Inzai, Chiba 270-1695, Japan

³ Department of Medical Technology, Faculty of Health Science, Gunma Paz College, Takasaki 370-0006, Japan



* Correspondence: hal@mail.pharm.tohoku.ac.jp; Tel.: +81-22-795-6824

Table of Contents

S2 ~ S4 NMR spectra of mucoroidiol (1)



S5 ~ S7 NMR spectra of firmibasiol (2)


¹H NMR (600 MHz, CDCl₃)




DEPT

¹³C NMR (150 MHz, CDCl₃)







DQFCOSY





HMQC spectrum







HMBC spectrum



NOESY spectrum

¹H NMR (600 MHz, CDCl₃)



DEPT

