

SUPPLEMENTARY TABLE S1. COMPARISON OF BASELINE CHARACTERISTICS OF INFANTS SEEN AND NOT SEEN AT VISIT 1 (12–15 MONTHS CORRECTED AGE)

	<i>Subjects seen (n=51)</i>	<i>Subjects not seen (n=110)</i>	<i>p</i>
Gestational age at birth (weeks) ^a	27.8±2.6	27.5±2.1	0.440
Male, <i>n</i> (%)	30 (59)	57 (52)	0.497
Race, white/black/Hispanic/other, <i>n</i> ^b	15/18/14/4	31/35/29/11	0.417
Birth			
Weight (g) ^a	893±204	921±197	0.404
Length (cm) ^a	34.1±3.1	34.6±2.7	0.308
HC (cm) ^a	24.2±2.0	24.4±1.8	0.657
Discharge			
Weight (g) ^a	2,753±871	2,946±1,076	0.264
Length (cm) ^a	46.1±3.9	46.9±4.9	0.329
HC (cm) ^a	33.0±2.4	33.5±3.0	0.305
Growth rate, birth to discharge			
Weight (g/day) ^a	21.0±3.4	22.0±3.6	0.091
Length (cm/week) ^a	0.96±0.21	0.96±0.23	0.892
HC (cm/week) ^a	0.70±0.15	0.72±0.15	0.442
Growth restriction (<10th percentile), <i>n</i> (%) ^c			
At birth	13 (25)	23 (21)	0.547
At discharge	28 (55)	50 (46)	0.312
Bronchopulmonary dysplasia, <i>n</i> (%)			
Moderate/severe	19 (37)	53 (48)	0.234
Intraventricular hemorrhage, <i>n</i> (%)			
Any	9 (18)	26 (24)	0.421
Grade 3/4	1 (2)	9 (8)	0.172
Late onset sepsis, <i>n</i> (%)	8 (16)	10 (9)	0.283
NEC, <i>n</i> (%)			
Medical	0 (0)	3 (3)	0.552
Surgical	0 (0)	2 (2)	0.564
Spontaneous intestinal perforation, <i>n</i> (%)	1 (2)	2 (2)	1.000
Calories at discharge [kcal/(kg·d)]	122±12	121±14	0.656
Receiving HM feeds at discharge, <i>n</i> (%)	28 (55)	39 (35)	0.026
PMA at discharge (weeks)	40.4±4.9	40.7±7.4	0.772
Length of stay (days)	88±43	93±56	0.576

^aMean ± standard deviation.

^bData not available for four subjects.

^cBased on Fenton 2013 growth curve.^{S1}

HC, head circumference; HM, human milk; NEC, necrotizing enterocolitis; NICU, neonatal intensive care unit; PMA, postmenstrual age.

SUPPLEMENTARY TABLE S2. COMPARISON OF BASELINE CHARACTERISTICS OF INFANTS SEEN AND NOT SEEN AT VISIT 2 (18–22 MONTHS CORRECTED AGE)

	Subjects seen (n=44)	Subjects not seen (n=117)	p
Gestational age at birth (weeks) ^a	27.9±2.6	27.5±2.1	0.308
Male, n (%)	25 (57)	62 (53)	0.724
Race, white/black/Hispanic/other ^b	14/15/11/4	32/38/32/11	0.212
Birth			
Weight (g) ^a	891±206	921±197	0.397
Length (cm) ^a	34.1±3.1	34.5±2.8	0.471
HC (cm) ^a	24.1±2.0	24.4±1.8	0.485
Discharge			
Weight (g) ^a	2,773±907	2,927±1,056	0.392
Length (cm) ^a	46.3±3.9	46.8±4.8	0.517
HC (cm) ^a	33.0±2.5	33.4±2.9	0.355
Growth rate, birth to discharge			
Weight (g/day) ^a	21.1±3.5	21.9±3.5	0.173
Length (cm/week) ^a	0.96±0.21	0.96±0.23	0.968
HC (cm/week) ^a	0.70±0.16	0.72±0.15	0.446
Growth restriction (<10th percentile), n (%) ^c			
At birth	12 (27)	24 (21)	0.793
At discharge	25 (57)	53 (46)	0.221
Bronchopulmonary dysplasia, n (%)			
Moderate/severe	17 (39)	55 (47)	0.378
Intraventricular hemorrhage, n (%)			
Any	8 (18)	27 (23)	0.532
Grade 3/4	1 (2)	9 (8)	0.288
Late onset sepsis, n (%)	8 (18)	10 (9)	0.099
NEC, n (%)			
Medical	0 (0)	3 (3)	0.563
Surgical	0 (0)	2 (2)	0.600
Spontaneous intestinal perforation, n (%)	1 (2)	2 (2)	1.000
Calories at discharge [kcal/(kg·d)] ^a	122±12	121±14	0.769
Receiving HM feeds at discharge, n (%)	25 (57)	42 (36)	0.020
PMA at discharge (weeks) ^a	40.6±5.2	40.6±7.2	0.992
Length of stay (days) ^a	89±45	92±55	0.706

^aMean ± standard deviation.

^bData not available for four subjects.

^cBased on the Fenton 2013 growth curve.^{S1}

HC, head circumference; HM, human milk; NEC, necrotizing enterocolitis; NICU, neonatal intensive care unit; PMA, postmenstrual age.

Supplementary Reference

- S1. Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr* 2013;13:59.