

**Alternative splicing of flowering time gene *FT* is associated with halving of time to flowering
in coconut**

Wei Xia^{2*}, Rui Liu^{1*}, Jun Zhang^{1*}, Annaliese S. Mason³, Zhiying Li¹, Shufang Gong¹, Yazhu Zhong¹,
Yajing Dou¹, Xiwei Sun¹, Haikuo Fan^{1&}, Yong Xiao^{1&}

¹ Coconut Research Institute, Chinese Academy of Tropical Agricultural sciences, Wenchang, P.R.
China

² College of tropical crops, Hainan University, Haikou, P.R. China

³ Department of Plant Breeding, IFZ Research Centre for Biosystems, Land Use and Nutrition, Justus
Liebig University Giessen, Heinrich-Buff-Ring 26-32, 35392i Giessen, Germany

>CCG011565-dwarf-CDS

ATGGCTAGCTGCGTGGATCCCCTTGTGGTAGGGAGGGTCATAGGCGAGGTGGTGGACCTTTTTGTTCCAC
CATGAGCATGTCTGTGAGCTACGGAACGAAGCATGTCAACAACGGCTGTGACGTGAAGCCGTCCATGGCTA
TCAACCCTCCAAGTGTCCAGATCGCAGGCCGACGAGCCGACCTCTATACTCTGGTGATGACCGATCCGGATG
CCCCAGTCCTAGTGACCCCACTATGAGGGAGTGGCTCCACTGGGTGGTGGTTAATATACCAGGTGGAACA
GATACCTCCAAG-----GAGAGGAGGTGGTGGCGTACATGGGACCGCGGCCCGCGGTGGGGATCCACCGC
TATGTCCTGGTGCTGTTCCAGCAGAAGTCAAGGTTCTTAACGGGGGCGGGGGTGGCTCTGCCTGCGTCTCG
TGCCAACTTCAACACCCGCACCTTCGCCGCCCGGCATGACCTTGGTCTCCCCGTCGCTGCGGTCTACTTCAAT
GCCCAGAAGGAGCCCGCCAACCGCCCGGTGA

> CCG011565-tall-CDS

ATGGCTAGCTGCGTGGATCCCCTTGTGGTAGGGAGGGTCATAGGCGAGGTGGTGGACCTTTTTGTTCCAC
CATGAGCATGTCTGTGAGCTACGGAACGAAGCATGTCAACAACGGCTGTGACGTGAAGCCGTCCATGGCTA
TCAACCCTCCAAGTGTCCAGATCGCAGGCCGACGAGCCGACCTCTATACTCTGGTGATGACCGATCCGGATG
CCCCAGTCCTAGTGACCCCACTATGAGGGAGTGGCTCCACTGGGTGGTGGTTAATATACCAGGTGGAACA
GATACCTCCAAGCGGCAGGAGAGGTGGTGGCGTACATGGGACCGCGGCCCGCGGTGGGGATCCAC
CGCTATGTCCTGGTGCTGTTCCAGCAGAAGTCAAGGTTCTTAACGGGGGCGGGGGTGGCTCTGCCTGCGTC
TCGTGCCAACTTCAACACCCGCACCTTCGCCGCCCGGCATGACCTTGGTCTCCCCGTCGCTGCGGTCTACTTC
AATGCCCAGAAGGAGCCCGCCAACCGCCCGGGAA

Supplementary file 2 Two transcripts sequences of *FT* gene (CCG011565.1)