

Supplementary Table 6. Diagnostic performances of LR-5 of LI-RADS 2018 and definite HCC of KLCA-NCC 2018 for diagnosing HCC on ECA-MRI and HBA-MRI in the subgroup with hepatitis B

	LR-5 of LI-RADS 2018 (95% CI)	Definite HCC of KLCA-NCC 2018 (95% CI)	P-value*
ECA-MRI (n=71)			
Sensitivity	75.0 (63.7–86.3)	67.9 (55.6–80.1)	0.094
Specificity	93.3 (80.7–100.0)	93.3 (80.7–100.0)	0.999
HBA-MRI (n=246)			
Sensitivity	68.5 (61.8–75.1)	79.1 (73.3–85.0)	<0.001
Specificity	96.6 (92.0–100.0)	94.9 (89.3–100.0)	0.313

LI-RADS, Liver Imaging Reporting and Data System; HCC, hepatocellular carcinoma; KLCA-NCC, Korean Liver Cancer Association-National Cancer Center; ECA, extracellular contrast agent; MRI, magnetic resonance imaging; HBA, hepatobiliary agent; CI, confidence interval.

*P-values between LR-5 of LI-RADS 2018 and definite HCC of KLCA-NCC 2018 by using the generalized estimating equation method.