

Supplementary Table 7. Diagnostic performances of LR-5/4 of LI-RADS 2018 and definite/probable HCC of KLCA-NCC 2018 for diagnosing HCC on ECA-MRI and HBA-MRI in the subgroup with hepatitis B

	LR-5/4 of LI-RADS 2018 (95% CI)	Definite/probable HCC of KLCA-NCC 2018 (95% CI)	P-value*
ECA-MRI (n=71)			
Sensitivity	87.5 (78.8–96.2)	83.9 (74.3–93.6)	0.313
Specificity	86.7 (69.5–100.0)	86.7 (69.5–100.0)	0.999
HBA-MRI (n=246)			
Sensitivity	87.7 (83.0–92.4)	85.0 (79.9–90.1)	0.056
Specificity	81.4 (71.4–91.3)	93.2 (86.8–99.6)	0.005

LI-RADS, Liver Imaging Reporting and Data System; HCC, hepatocellular carcinoma; KLCA-NCC, Korean Liver Cancer Association-National Cancer Center; ECA, extracellular contrast agent; MRI, magnetic resonance imaging; HBA, hepatobiliary agent; CI, confidence interval.

*P-values between LR-5/4 of LI-RADS 2018 and definite/probable HCC of KLCA-NCC 2018 by using the generalized estimating equation method.