

Supplementary Table 2  
*Outcomes organized by methodologically relevant variables*

	Q	k	Hedge's <i>g</i>	SE	95% CI	
					LL	UL
<b>Comparison</b>	0.04					
Neutral cue		55	0.77***	0.07	0.64	0.91
Baseline		13	0.75***	0.14	0.48	1.02
<b>Cue Presentation Order</b>	0.16					
Other		37	0.82***	0.09	0.65	0.99
Fixed		14	0.75***	0.14	0.48	1.03
<b>Craving Measure</b>	9.27					
Single item		24	0.69***	0.10	0.50	0.88
QSU-10		23	0.87***	0.10	0.68	1.07
QSU-4		11	0.76***	0.14	0.48	1.04
Cigarette Craving Questionnaire		7	0.96***	0.18	0.62	1.31
Shiffman-Jarvik		2	-0.06	0.39	-0.82	0.70
QSU Factor 1		1	0.19	0.44	-0.68	1.06
<b>Abstinence Requirement</b>	5.29*					
Any abstinence		40	0.66***	0.07	0.52	0.80
No abstinence		28	0.92***	0.09	0.75	1.09
<b>Cue Presentation Mode</b>	15.25**					
<i>In vivo</i>		26	0.62***	0.09	0.46	0.79
Images		24	0.80***	0.10	0.61	0.98
Scripted imagery		13	1.11***	0.12	0.87	1.35
Virtual reality		3	0.78**	0.27	0.25	1.31
Video		2	0.09	0.31	-0.53	0.70

Note. Q = Cochran's *Q*; k = number of studies; SE = standard error; CI = confidence interval; LL = lower limit; UL = upper limit.

\*p < .05, \*\*p < .01, \*\*\*p < .001