

***Mycobacterium tuberculosis* resistance prediction and lineage classification
from genome sequencing: comparison of automated analysis tools**

Viola Schleusener¹, Claudio U. Köser², Patrick Beckert^{1,3}, Stefan Niemann^{1,3*}, and Silke Feuerriegel^{1,3*}

¹ Division of Molecular and Experimental Mycobacteriology Group, Research Center Borstel, Borstel, Germany

² Department of Medicine, University of Cambridge, Cambridge, United Kingdom

³ German Center for Infection Research, Borstel Site, Borstel, Germany


Corresponding author: Stefan Niemann, Division of Molecular and Experimental Mycobacteriology Group, Research Center Borstel, Borstel, Germany. Tel: 0049-4537-1887620, E-Mail: sniemann@fz-borstel.de

*SN and SF contributed equally to this work


Supplementary Figure S1.

Mapping results (stampy) for Mykrobe Predictor TB's low-frequency variants in *rpoB* (n=7) and *katG* (n=1)


IGV mapping of 1327-04 at genome position 2155168




IGV mapping of 3158-04 at genome position 761154




IGV mapping of 4712-04 at genome position 761079




IGV mapping of 4717-04 at genome position 761139




IGV mapping of 6640-04 at genome position 761154



IGV mapping of 8082-03 at genome position 761133



IGV mapping of 10205-03 at genome position 761133



IGV mapping of 11818-03 at genome position 761108

