

Supplementary Materials

TITLE: Low serum lymphocyte level is associated with poor exercise capacity and quality of life in chronic obstructive pulmonary disease

Sung Woo Moon¹, Ah Young Leem¹, Young Sam Kim¹, Ji-Hyun Lee², Tae-Hyung Kim³, Yeon-Mok Oh⁴, Hyejung Shin⁵, Joon Chang¹, Ji Ye Jung¹, and KOLD Study Group

Affiliations:

¹Division of Pulmonology, Department of Internal Medicine, Severance Hospital, Yonsei University College of Medicine., Seoul, Republic of Korea

²Department of Internal Medicine, CHA Bundang Medical Center, CHA University, Seongnam, Republic of Korea

³Division of Pulmonary and Critical Care Medicine, Hanyang University Guri Hospital, Hanyang University College of Medicine, Guri, Republic of Korea

⁴Department of Pulmonary and Critical Care Medicine and Clinical Research Center for Chronic Obstructive Airway Diseases, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea

⁵Biostatistics Collaboration Unit, Department of Biomedical Systems Informatics, Yonsei University College of Medicine, Seoul, Korea

Supplementary Figure 1. Serum lymphocyte percent (%) over the follow-up period.

