

SUPPLEMENTARY TABLES:

Supplementary Table 1: Chest imaging for three groups

	COVID-19 alone (n=108)	Cirrhosis+COVID-19 (n=37)	Cirrhosis alone (n=127)	P-value for all three groups
Chest X ray (done in everyone)				
Normal	9	4	32	<0.0001
Bilateral Ground glass opacities	33	11	7	<0.0001
Bilateral Ground glass consolidation	5	1	2	0.38
Pneumonia	46	11	11	<0.0001
Pulmonary Edema	14	4	8	0.21
Pleural effusion	12	6	19	0.61
CT Scan	COVID-19 alone (n=57)	Cirrhosis+COVID-19 (n=21)	Cirrhosis alone (n=56)	
Normal	0	0	2	
Bilateral Ground glass opacities	23	7	9	0.006
Bilateral Ground glass consolidation	6	2	2	0.23
Pneumonia	19	6	7	0.01
Pulmonary Edema	2	3	4	0.25
Pleural effusion	6	2	15	0.18

Data presented as raw numbers; Kruskal-Wallis, Chi-square tests or ANOVA as appropriate. There could be more than one finding per imaging, which is why the numbers do not add up to the total above.

Supplementary Table 2: Medications used for COVID-19 patients

	COVID-19 alone (n=108)	COVID-19 +Cirrhosis (n=37)
Antibiotics		
Beta lactams	45	11
Fluoro quinolones	3	2
Zosyn/Timentin/Augmentin	7	4
Clindamycin	3	1
Vancomycin	14	9
Daptomycin	0	0
Macrolides	22	10
Tetracyclines	23	2
Anti-fungal	4	1
Imipenem/Meropenem/Ertapenem	4	0
Metronidazole	3	0
Linezolid	1	0
Others	5	6
Non-antibiotic medications		
NSAIDs	5	0
Acetaminophen	65	19
Chloroquine/Hydroxychloroquine	46	11
Tocilizumab	12	9
Lopinavir/Ritonavir	1	0
Remdesivir	2	1
Sarilumab	1	1
Clinical Trials		
Enrolled in Clinical trial*	46	8

Data presented as raw numbers; *:p<0.05 between the 2 groups ; none of the other comparisons were significant. Comparisons were performed using Chi-square tests.