

Baseline Characteristics from UNITE: An Observational, International, Multicentre Registry to Evaluate Hidradenitis Suppurativa in Clinical Practice

Authors and Affiliations:

Errol P. Prens, MD, PhD¹; Aida M Lugo-Somolinos, MD²; Amy S. Paller, MD, FAAD³; Francisco Kerdel, MD⁴; Yinghui Duan, PhD⁵; Henrique D. Teixeira, PhD, MBA⁵; Michelle Longcore, PharmD⁵; Alexa B. Kimball, MD, MPH⁶

1 Erasmus University Medical Center, Rotterdam, Netherlands; or CHU de Reims, Hôpital Robert Debré, Service de Dermatologie, Reims Cedex, France

2 University of North Carolina Hospital, Dermatology and Skin Cancer Center, Chapel Hill, NC, USA

3 Northwestern University Feinberg School of Medicine, Chicago, IL, USA

4 Florida Academic Dermatology Centers, Miami, FL, USA

5 AbbVie, Inc, North Chicago, IL, USA

6 Harvard Medical School and Beth Israel Deaconess Hospital, Boston, MA, United States

Journal: American Journal of Clinical Dermatology

Counts: Words: Abstract 239; Body 3001. Tables:7. Figures: 1. Online Resource: 1

Corresponding author:

Errol P Prens, MD, PhD
Erasmus University Medical Center
Dr. Molewaterplein 40
Post Box 2040
Rotterdam 3014 GD, Netherlands
Phone: (31) 10 703 4580
Email: e.prens@erasmusmc.nl

Online Resource 1. Lesion Assessment

Body Area	Anatomic Region	Body Area	Anatomic Region
Mammary	Intermammary area	Other	Scalp
	Sub/inframammary area, left		Face ^a
	Sub/inframammary area, right		Neck
Pelvic	Buttock, left		Abdomen
	Buttock, right		Chest ^b
	Inguino-cural fold, left		Back
	Inguino-cural fold, right		Pubic region
	Perianal		Genital region
	Perineal		Upper extremity, left
Axilla	Axilla, left		Upper extremity, right
	Axilla right		Lower extremity, left
		Lower extremity, right	

a. includes ears and area behind the ears. b. excludes the mammary, inframammary, intermammary areas.