

Exogenous abscisic acid induces the lipid and flavonoid metabolism of tea plants under drought stress

Zhongshuai Gai^{12#}, Yu Wang^{1#}, Yiqian Ding¹, Wenjun Qian¹, Chen Qiu¹, Hui Xie¹, Litao Sun¹, Zhongwu Jiang², Qingping Ma³, Linjun Wang⁴, Zhaotang Ding^{1*}

¹Tea Research Institute, Qingdao Agricultural University, Qingdao 266109, China

²College of Life Science, Yantai University, Yantai, Shandong, 264005, China

³College of agriculture, Liaocheng University, Liaocheng, Shandong, 252059, China

⁴Fruit tea station of weihai agricultural and rural affairs service center, Weihai, Shandong, 264200, China

These authors contributed equally to this study.

*CORRESPONDENCE: Dr Zhaotang Ding E-mails: dzttea@163.com

Table S1 Quality of sequencing

Samples	Clean reads	Cleans bases			GC(%)	
		(G)	Error (%)	Q20 (%)		
CK1	53,324,234	8.0	0.34	97.29	92.97	44.52
CK2	50,912,316	7.64	0.36	97.09	92.54	44.50
CK3	49,215,718	7.38	0.33	97.37	93.22	44.70
MD1	47,871,864	7.18	0.33	97.34	93.18	45.06
MD2	53,180,726	7.98	0.32	97.48	93.44	44.80
MD3	52,383,110	7.86	0.33	97.38	93.34	44.98
AT1	51,863,816	7.78	0.33	97.35	93.29	45.14
AT2	53,845,492	8.08	0.33	97.37	93.41	45.61
AT3	52,340,128	7.85	0.32	97.48	93.66	45.66
SD1	51,611,966	7.74	0.33	97.35	93.20	44.89
SD2	51,740,376	7.76	0.33	97.40	93.36	45.01
SD3	51,528,212	7.73	0.34	97.27	93.10	44.57

CK: control group; MD: mild drought group; AT: ABA treatment group; SD: severe drought group