## **INTESTINAL RESEARCH**

	IST	+	IST	-		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% Cl	M–H, Fixed, 95% Cl
Abreu 2016	16	148	4	38	11.4%	1.03 [0.32, 3.28]	
Al Taweel 2016	1	99	6	47	16.1%	0.07 [0.01, 0.60]	←
Andrisani 2013	10	70	3	22	7.8%	1.06 [0.26, 4.24]	
Cem Çekiç 2015	19	47	15	26	23.1%	0.50 [0.19, 1.32]	
Mantzaris 2015	7	28	9	47	10.1%	1.41 [0.46, 4.32]	•
Ministro 2011	2	14	3	10	6.0%	0.39 [0.05, 2.92]	
Papay 2011	7	131	8	59	20.9%	0.36 [0.12, 1.04]	
Ramos 2013	0	20	1	5	4.6%	0.07 [0.00, 2.11]	· · · · · · · · · · · · · · · · · · ·
Total (95% CI)		557		254	100.0%	0.57 [0.37, 0.89]	•
Total events	62		49				
Heterogeneity: $\text{Chi}^2 = 10.29$ , df = 7 ( <i>P</i> =0.17); $l^2 = 32\%$							
Test for overall effect: $Z = 2.48$ ( $P = 0.01$ )							0.01 0.1 1 10 100 Favours [experimental] Favours [control]

**Supplementary Fig. 1.** OR positive interferon-gamma release assays in immunosuppressed patients (IST+) compared to non-immunosuppressed (IST-) with fixed model. IST, immunosuppressive therapy; M-H, Mantel-Haenszel method.