


SUPPLEMENTARY INFORMATION FOR

# Explainable artificial intelligence model to predict acute critical illness from electronic health records

**Simon Meyer Lauritsen<sup>1,2</sup>, Mads Kristensen<sup>1</sup>, Mathias Vassard Olsen<sup>3</sup>, Morten Skaarup Larsen<sup>3</sup>,  
Katrine Meyer Lauritsen<sup>2</sup>, Marianne Johansson Jørgensen<sup>4</sup>,  
Jeppe Lange<sup>2,4</sup>, Bo Thiesson<sup>1,5</sup>**

<sup>1</sup>*Enversion A/S, Fiskerivej 12, 1st floor, 8000 Aarhus C, Denmark;* <sup>2</sup>*Department of Clinical Medicine,  
Aarhus University, Palle Juul-Jensens Boulevard 82, 8200 Aarhus N, Denmark;* <sup>3</sup>*Department of  
Biomedical Engineering and Informatics, Aalborg University, Niels Jernes Vej 12, 9220 Aalborg Ø,  
Denmark;* <sup>4</sup>*Department of Research, Horsens Regional Hospital, Sundvej 30, 8700 Horsens,  
Denmark;* <sup>5</sup>*Department of Engineering, Aarhus University, Inge Lehmanns Gade 10, 8000 Aarhus C,  
Denmark.*

\*Corresponding author: Simon Meyer Lauritsen (sla@enversion.dk)



**Supplementary Figure 1 | Cross validation scheme.** Data were randomly divided into 5 portions of 20% each. For each fold four portions (80 %) was used to fit the xAI-EWS model parameters during training. The remaining 20% was split into two portions of 10% each for validation and test. The validation data were used to perform an unbiased evaluation of a model fit during training, and the test data were used to provide an unbiased evaluation of the final model. For each fold data were shifted such that a new portion was used for testing. All data for a single patient was assigned to either train, validation or test data.

**Supplementary Table 1 | The Area Under The Receiver Operating Characteristics (AUROC) curve**

Hours before onset	Fold	Sepsis				Acute Kidney Injury				Acute Lunge Injury			
		TCN	SOFA	MEWS	GB-Vital	TCN	SOFA	MEWS	GB-Vital	TCN	SOFA	MEWS	GB-Vital
0	1	0.92	0.83	0.8	0.84	0.88	0.75	0.70	0.76	0.90	0.80	0.85	0.65
0	2	0.95	0.84	0.82	0.83	0.87	0.74	0.68	0.78	0.90	0.80	0.89	0.61
0	3	0.92	0.83	0.82	0.85	0.88	0.74	0.72	0.78	0.93	0.77	0.85	0.63
0	4	0.90	0.83	0.83	0.85	0.9	0.75	0.71	0.76	0.9	0.77	0.81	0.65
0	5	0.93	0.83	0.84	0.83	0.86	0.73	0.70	0.76	0.89	0.82	0.84	0.63
3	1	0.88	0.80	0.78	0.81	0.85	0.74	0.68	0.70	0.85	0.74	0.85	0.63
3	2	0.85	0.80	0.79	0.81	0.84	0.73	0.68	0.72	0.88	0.74	0.78	0.62
3	3	0.85	0.79	0.78	0.81	0.85	0.74	0.70	0.72	0.88	0.72	0.78	0.61
3	4	0.89	0.79	0.78	0.80	0.83	0.75	0.68	0.71	0.85	0.73	0.81	0.55
3	5	0.87	0.8	0.80	0.81	0.86	0.73	0.68	0.75	0.89	0.75	0.85	0.63
6	1	0.86	0.78	0.78	0.78	0.84	0.74	0.70	0.70	0.83	0.73	0.81	0.61
6	2	0.85	0.77	0.77	0.77	0.85	0.72	0.68	0.69	0.82	0.69	0.78	0.63
6	3	0.86	0.79	0.77	0.78	0.85	0.74	0.70	0.69	0.84	0.72	0.78	0.60
6	4	0.86	0.76	0.78	0.80	0.81	0.74	0.69	0.70	0.85	0.75	0.78	0.59
6	5	0.83	0.79	0.77	0.80	0.83	0.73	0.67	0.68	0.88	0.70	0.71	0.63
12	1	0.84	0.77	0.68	0.78	0.85	0.72	0.61	0.68	0.82	0.70	0.74	0.60
12	2	0.83	0.76	0.71	0.77	0.85	0.71	0.69	0.68	0.86	0.76	0.75	0.59
12	3	0.84	0.77	0.71	0.78	0.86	0.72	0.66	0.65	0.84	0.76	0.83	0.61
12	4	0.85	0.77	0.72	0.79	0.85	0.72	0.67	0.68	0.85	0.72	0.77	0.61
12	5	0.84	0.75	0.72	0.79	0.83	0.71	0.61	0.65	0.82	0.70	0.80	0.61
24	1	0.81	0.72	0.71	0.79	0.8	0.66	0.62	0.68	0.85	0.67	0.70	0.60
24	2	0.81	0.72	0.71	0.79	0.77	0.66	0.61	0.60	0.84	0.65	0.71	0.60
24	3	0.80	0.72	0.67	0.77	0.77	0.62	0.60	0.65	0.82	0.70	0.74	0.59
24	4	0.82	0.65	0.65	0.77	0.81	0.67	0.62	0.63	0.85	0.67	0.76	0.58
24	5	0.77	0.71	0.61	0.76	0.79	0.62	0.62	0.60	0.83	0.67	0.75	0.58

TCN: Temporal Convolutional Network, SOFA: Sequential Organ Failure Assessment, MEWS: Early Warning Score, GB-Vital: Gradient Boosting model based on vital sign parameters.

**Supplementary Table 2 | Area Under The Precision Recall (AUPRC) Curve**

Hours before onset	Fold	Sepsis				Acute Kidney Injury				Acute Lung Injury			
		TCN	SOFA	MEWS	GB-Vital	TCN	SOFA	MEWS	GB-Vital	TCN	SOFA	MEWS	GB-Vital
0	1	0.50	0.12	0.12	0.42	0.24	0.11	0.07	0.18	0.24	0.05	0.24	0.05
0	2	0.48	0.11	0.14	0.42	0.21	0.14	0.07	0.20	0.25	0.05	0.12	0.05
0	3	0.38	0.09	0.14	0.42	0.21	0.12	0.06	0.18	0.21	0.04	0.24	0.07
0	4	0.39	0.12	0.13	0.45	0.19	0.12	0.05	0.20	0.25	0.05	0.18	0.05
0	5	0.40	0.11	0.13	0.41	0.23	0.12	0.07	0.17	0.22	0.05	0.19	0.07
3	1	0.12	0.07	0.09	0.10	0.19	0.12	0.08	0.08	0.22	0.05	0.15	0.05
3	2	0.22	0.07	0.06	0.09	0.19	0.11	0.08	0.08	0.23	0.04	0.10	0.04
3	3	0.16	0.09	0.09	0.14	0.23	0.12	0.06	0.07	0.18	0.06	0.20	0.04
3	4	0.18	0.06	0.12	0.11	0.22	0.10	0.05	0.07	0.25	0.05	0.10	0.05
3	5	0.24	0.07	0.08	0.13	0.22	0.10	0.07	0.07	0.22	0.05	0.12	0.05
6	1	0.15	0.06	0.07	0.08	0.21	0.10	0.07	0.03	0.23	0.05	0.15	0.05
6	2	0.12	0.06	0.06	0.05	0.21	0.09	0.05	0.02	0.25	0.03	0.10	0.05
6	3	0.15	0.05	0.12	0.05	0.20	0.11	0.08	0.05	0.19	0.04	0.20	0.07
6	4	0.15	0.05	0.07	0.07	0.19	0.09	0.07	0.02	0.26	0.05	0.09	0.03
6	5	0.13	0.06	0.09	0.05	0.21	0.10	0.06	0.03	0.18	0.05	0.12	0.05
12	1	0.11	0.05	0.04	0.05	0.19	0.08	0.07	0.03	0.20	0.02	0.05	0.05
12	2	0.10	0.05	0.07	0.04	0.19	0.10	0.06	0.02	0.24	0.02	0.04	0.05
12	3	0.12	0.04	0.07	0.05	0.20	0.07	0.06	0.02	0.20	0.05	0.08	0.03
12	4	0.12	0.05	0.07	0.04	0.21	0.08	0.07	0.02	0.21	0.05	0.07	0.03
12	5	0.11	0.09	0.05	0.07	0.19	0.08	0.05	0.03	0.22	0.02	0.07	0.05
24	1	0.09	0.07	0.04	0.07	0.13	0.04	0.02	0.04	0.24	0.02	0.04	0.05
24	2	0.07	0.06	0.05	0.05	0.14	0.05	0.02	0.02	0.22	0.02	0.04	0.04
24	3	0.08	0.04	0.05	0.05	0.13	0.03	0.03	0.02	0.22	0.02	0.05	0.02
24	4	0.08	0.05	0.05	0.08	0.12	0.04	0.04	0.02	0.24	0.02	0.06	0.03
24	5	0.08	0.08	0.03	0.08	0.13	0.04	0.02	0.02	0.23	0.05	0.02	0.02

TCN: Temporal Convolutional Network, SOFA: Sequential Organ Failure Assessment, MEWS: Early Warning Score, GB-Vital: Gradient Boosting model based on vital sign parameters.

