

Supplementary Table 4. Multivariable analysis of BMI stratified by treatment group

Response	OR	95% CI	p-value
<i>Total Cohort</i>			
Overweight vs Normal	0.84	0.45-1.60	0.60
Obese vs Normal	0.58	0.31-1.09	0.09
<i>Combination Therapy</i>			
Overweight vs Normal	1.04	0.25-4.30	0.95
Obese vs Normal	0.47	0.12-1.84	0.28
<i>Monotherapy</i>			
Overweight vs Normal	0.79	0.38-1.66	0.54
Obese vs Normal	0.64	0.31-1.32	0.23
PFS	HR	95% CI	p-value
<i>Total Cohort</i>			
Overweight vs Normal	1.04	0.71-1.52	0.84
Obese vs Normal	1.28	0.90-1.83	0.18
<i>Combination Therapy</i>			
Overweight vs Normal	1.31	0.53-3.19	0.56
Obese vs Normal	1.64	0.69-3.91	0.26
<i>Monotherapy</i>			
Overweight vs Normal	0.98	0.64-1.50	0.93
Obese vs Normal	1.20	0.81-1.79	0.36
OS	HR	95% CI	p-value
<i>Total Cohort</i>			
Overweight vs Normal	1.10	0.71-1.70	0.67
Obese vs Normal	1.10	0.72-1.67	0.65
<i>Combination Therapy</i>			
Overweight vs Normal	0.91	0.30-2.74	0.86
Obese vs Normal	1.31	0.50-3.43	0.58
<i>Monotherapy</i>			
Overweight vs Normal	1.14	0.70-1.86	0.59
Obese vs Normal	1.05	0.55-1.68	0.85

Adjusted for age, stage, and prior therapy