

# THE LANCET

## Global Health

### Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed.  
We post it as supplied by the authors.

Supplement to: Busch-Hallen J, Walters D, Rowe S, Chowdhury A, Arabi M. Impact of COVID-19 on maternal and child health. *Lancet Glob Health* 2020; published online Aug 1. [http://dx.doi.org/10.1016/S2214-109X\(20\)30327-2](http://dx.doi.org/10.1016/S2214-109X(20)30327-2).

Table 1. Estimated number of deaths of children year attributable to reductions in breastfeeding prevalence by region and country income group during the Covid-19 pandemic. Child deaths are due to pneumonia and diarrhea .

	<b>Scenario 1</b>	<b>Scenario 2</b>	<b>Scenario 3</b>	<b>Scenario 4</b>
	5% Relative Reduction in BF Prevalence	10% Relative Reduction in BF Prevalence	25% Relative Reduction in BF Prevalence	50% Relative Reduction in BF Prevalence
<b>Regions</b>	<b>Total child deaths</b>			
<b>East Asia &amp; Pacific</b>	970	1,798	4,112	7,488
<b>Europe &amp; Central Asia</b>	87	171	413	1,719
<b>Middle East &amp; North Africa</b>	223	440	1,059	1,995
<b>Latin America &amp; Caribbean</b>	194	383	917	1,719
<b>South Asia</b>	7,035	13,750	32,187	58,226
<b>Sub-Saharan Africa</b>	7,960	15,596	36,766	67,252
<b>Total</b>	<b>16,469</b>	<b>32,139</b>	<b>75,455</b>	<b>138,398</b>