Table S1b: P-values and significance levels for all comparisons of retinal thickness measurement means derived from SD-OCT. Multiple comparisons of means with Tukey's approach were performed between different measurement positions at 1, 2, 3, 4, and 6mm ventral to the central and nasal ONH rim. Comparison between positions x and y is indicated as x - y. Asterisks indicate significance levels for all significant differences: (\*)  $P \le 0.05$ , (\*\*)  $P \le 0.01$  and (\*\*\*)  $P \le 0.001$ .

P-values and significance levels for all comparisons of SD-OCT-derived thickness measurements											
Comparisons between distances from ONH		1-2	1-3	1 - 4	1 - 6	2 - 3	2 - 4	2 - 6	3 - 4	3 - 6	4 - 6
IR	nasal	***	***	0.99	***	***	***	***	***	***	***
	central	0.19	***	***	***	***	***	*	***	***	***
INL	nasal	***	***	***	***	***	***	***	***	***	***
	central	***	***	***	***	***	***	***	0.95	***	***
ONL	nasal	***	***	***	***	***	*	***	0.13	***	***
	central	0.99	***	***	*	***	***	**	0.19	***	***
OR	nasal	***	***	***	**	***	***	***	***	***	***
	central	***	***	***	***	***	***	***	***	***	***
PR	nasal	***	***	***	***	***	***	***	***	***	***
	central	***	***	***	***	***	***	***	***	***	***