Study	Study question or objective clearly stated?	Population clearly and fully described, including a case definition?	Cases were consecutive?	Subjects were comparable?	Outcome measures clearly defined, valid, reliable, and implemented consistently across all study participants?	Length of follow-up?	Statistical methods well- described?	Results well- described?
Arentz et al, JAMA <sup>12</sup>	Y	Y	Y	Y	Y	5-13 days	n/a	Y
Brill et al, preprint <sup>13</sup>	Y	Y	Y	Y	Y	5-29 days	Y	Y
Cai et al., Eur J All & Clinical Imm, <sup>14</sup>	Y	Y	Y	Y	Y	29-55 days	Y	Y
Chan et al, preprint <sup>15</sup>	Y	Y	Y	Y	Y	Until discharged or deceased	Y	Y
Chen et al., BMJ <sup>16</sup>	Y	Y	Y	Y	Y	16-48 days	Y	Y
Chen et al, JCI <sup>17</sup>	Y	Y	Y	Y	Y	0-27 days	Y	Y
Cheng et al, KI <sup>18</sup>	Y	Y	Y	Y	Y	18-32 days	Y	Y
Guan et al, NEJM <sup>4</sup>	Y	Y	Y	Y	Y	2-51 days	Y	Y
Hirsch et al, KI <sup>19</sup>	Y	Y	Y	Y	Y	7-42 days	Y	Y
H ( 1 MAT20	V	N	V	V	V	Start date uncertain. Follow-up until March	Y	V
Hong et al, YMJ <sup>20</sup>	Y	N	Y	Y	Y	29	Y	Y
Huang et al, Lancet <sup>3</sup>	Y	Y	Y	Y	Y	20-37 days 49-73 days	Y	Y
Jiang et al, preprint <sup>21</sup>	Y	Y	Υ	Y	Y	49-73 days	Y	Y
Pei et al, JASN,								
May <sup>22</sup>	Y	Y	Y	Y	Y	14-26 days	Y	Y
Qiu et al, JMV <sup>23</sup>	Y	Y	Y	Y	Y	11-32 days	Y	Y
Regina et al, preprint <sup>24</sup>	Y	Y	Y	Y	Y	14-39 days	Y	Y
Richardson, JAMA <sup>25</sup>	Y	Y	Y	Y	Y	0-35 days	Ν	
Rubin et al.,								
preprint <sup>26</sup>	Y	Y	Y	Y	Y	0-42 days	Y	Y
Shi et al, JAMA <sup>27</sup>	Y	Y	Y	Y	Y	5-26 days	Y	Y
Shi et al, preprint <sup>28</sup> Wang et al, JAMA <sup>29</sup>	Y Y	Y Y	Y Y	Y Y	Y Y	Until discharged or deceased 28-34 days	Y Y	Y Y
Wang et al., Crit Care <sup>30</sup>	Y	Y	Y	Y	Y	Until discharged or deceased	Y	Y

Supplemental Table 1. Quality assessment of included studies based on the NHLBI Case Series Quality Assessment Tool.

Wang et al., J								
Infection <sup>31</sup>	Y	Y	Y	Y	Y	28-65 days	Y	Y
Xiao et al, preprint <sup>32</sup>	Y	Y	Y	Y	Y	14-62 days	Y	Y
Yan et al, preprint <sup>33</sup>	Y	Y	Y	Y	Y	0-51 days	Y	Y
Yang et al, Lancet <sup>34</sup>	Y	Y	Y	Y	Y	14-47 days	Y	Y
Zhang et al, Clin Micr						-		
Inf <sup>35</sup>	Υ	Y	Y	Y	Y	3-29 days	Y	Y
Zhao et al, preprint <sup>36</sup>	Y	Y	Y	Y	Y	21-39 days	Y	Y
Zheng et al, J								
Zheijang University <sup>37</sup>	Υ	Y	Y	Y	Y	0-43 days	Y	Y
						Until discharged or		
Zhou et al, Lancet <sup>38</sup>	Y	Y	Υ	Y	Υ	deceased	Y	Y
						Until discharged or		
Zhou et al, preprint <sup>39</sup>	Υ	Y	Y	Y	Y	deceased	Ν	Y



Section/topic	#	Checklist item	Reported on page #
TITLE	•		
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4, 5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5, 6
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	5
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5, 6
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6, 7
Risk of bias in individual studies	12	P Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7



## PRISMA 2009 Checklist

Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency	7	
		(e.g., l <sup>2</sup> ) for each meta-analysis.		

	•	Page 1 of 2			
Section/topic	#	Checklist item			
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	7		
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.			
RESULTS					
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	7		
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	7, 8		
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	8		
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	8, 9		
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	9		
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	10		
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).			
DISCUSSION					
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	10, 11		
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	12		
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	13		
FUNDING	1				
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	13		



## **PRISMA 2009 Checklist**

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Page 2 of 2